

President's Message

The 24th IAGS is now less than 12 months away and I urge members and other potential attendees, especially graduate students, to

David Cohen start putting pen to paper (well, finger to keyboard) on their abstracts. With the ongoing mineral exploration boom, AAG hopes to have a large contingent from industry attending the conference in New Brunswick, Canada next June. The Symposium technical program will provide a range of broad themes, from those in "classical" exploration geochemistry to environmental and other areas of applied geochemistry. The pre- and post-symposium field excursions should also attract new faces to the IAGS. Dave Lentz and the Local Organizing Committee have worked to keep costs as low as possible to encourage student participation, though delegates can opt for upmarket accommodation options if they want. Further details about the Symposium are provided in this edition of EXPLORE and the AAG website. With the Symposium in mind, I would like to draw member's attention to the new set of practical guidelines for the preparation and delivery of oral and poster papers by Ian Robertson (see AAG News on the AAG web site).

Apart from being inundated with general geochemical consulting work, word from a few of the AAG members who are general geochemical consultants is that there is almost as much demand for training courses for company personnel. This is training in the basics from geochemical processes in surface environments to sampling, analysis, QC and simple data analysis. A couple of short courses that I have recently run with Neil Rutherford in Beijing at CUG attracted a total of 120 students and staff.

The AAG already has a significant commitment to the educational side of the discipline. These include support



of the AAG Distinguished Lecturer (currently Kurt Kyser), the IAGS with its accompanying workshops, AAGsponsored short courses and workshops, some lectures and the like posted on the AAG website, support for students attending symposia and the donation of journal and book collections to universities in developing countries. There are many of our more senior members who may be losing some enthusiasm for spending long periods in the field battling black flies, black snakes and black rhinos, but who are both enthusiastic and well-prepared to transfer their knowledge and experience to the next generation within industry or those universities where applied geochemistry is lacking. There is certainly scope for the AAG to expand its educational role and to act as a contact point for companies and even universities seeking the services of AAG members in delivering customized educational programs. AAG could list members on the website with the training modules that they can offer (subjects, duration, etc) and develop a repository of teaching materials that could be drawn upon.

I would like to acknowledge the ongoing work of Eion Cameron handling the AAG investments (a major source of operating funds for the Association) during this period of financial market instability and stock market upheavals. We also note Gwendy Hall's retirement from official duties with the Geological Survey of Canada but look forward to her continuing activities in the geochemical sphere. I also need to make one correction to my last letter - the Distinguished Applied Geochemist Fund will be chaired by our immediate past-president Rob Bowell.

Dave Cohen, President

The geochemistry of acid-saline waters and drainage sediments in the Avon Catchment, Western Australian Wheatbelt

Newsletter for the Association of Applied Geochemists




Figure 1: Location of Australian Wheatbelt east of Perth in Western Australia. many parts of the West Australian (WA) Wheatbelt (Fig. 1). Engineering options to counter rising saline groundwaters are increasingly being adopted in the WA Wheatbelt (Clarke et al. 2002; Hatton et al. 2003).

Introduction

The relationship between native vegetation clearance and increasing groundwater and surface water salinities is well established in the semi-arid regions of Australia (Allison et al. 1990). The primary cause of dryland salinity in many parts of Australia is due to increased groundwater recharge following native vegetation clearance for annual crops and pastures. This has caused mobilisation of salt stored in the shallow regolith and/ or increased groundwater discharge. Salinisation due to rising water tables is a significant management issue in

continued on page 2













PAGE 2 NUMBER 140 EXPLORE

The geochemistry of acid-saline waters and drainage... continued from page 1

The adoption of engineered drainage solutions to control groundwater levels can be an effective management tool in landscapes affected by dryland salinity. However, hydrological efficiency is highly dependant on near surface regolith properties. Recently, there has been extensive state government agency and landholder interest in using deep (2-3 m) drainage systems as an option to control water table rise (Ali et al. 2004a) (Fig. 2). Over 10,000 kilometres of drains have already been constructed to protect both lowlying land from salinisation and rehabilitate marginally saline lands, and plans are already being assessed for further expansion of drainages on individual properties, and to establish regional integrated drainage systems.

Concerns have been raised by landholders, local government authorities, land management agencies and the general community (Dogramaci & Degens 2003) regarding the potential for drainage systems to discharge acidic, trace element rich waters that may impact on receiving environments. Initial investigations of some major drainage schemes found that these can discharge waters of pH 2-3 with total dissolved solids (TDS) of 30000 to 50000 mg l⁻¹ (TDS of seawater is 34600 mg l⁻¹) at 5-10 ML per day (Ali et al. 2004b).

In order to address these concerns, and provide a robust scientific basis upon which to plan drainage schemes and assess geochemical risks of acid discharge, the Engineering



Figure 2. Typical deep acidic saline drain with in the WA Wheatbelt.

Evaluation Initiative (EEI) and the Wheatbelt Drainage Evaluation Project have been commissioned as part of the WA National Action Plan for Salinity and Water Quality (NAPSWQ). Although salinity was originally assumed to be the main potential water quality problem, the realisation

continued on page 4

Notes from the Editor

The September issue of **EXPLORE** (No. 140) includes two articles. Paul Shand, Brad Degens, and Steve Rogers describe acid-saline waters in the Western Australian Wheatbelt. Dave Lentz provides an overview of the United Nations International Year of Planet Earth (2008). Scientific and technical editing assistance for this issue was provided by Matt Leybourne, GNS Science and Wendy Spirito, Geological Survey of Canada. Thanks to all authors and editors for their contributions. This issue also includes

TABLE OF CONTENTS

President's Message
The geochemistry of acid-saline waters and drainage sediments
in the Avon Catchment, Western Australian Wheatbelt1
Notes from the Editor2
The United Nations International Year of Planet Earth (2008)12
24th International Applied Geochemistry Symposium (IAGS 2009) 14
AAG Student Paper Competition17
Calendar of Events
Mineral Exploration Round Up Short Course19
Recent Papers
Modular Course in Exploration Geochemists24
Membership Application
GRSG Annual Meeting Call for Papers26
List of Advertisers


detailed information about the upcoming 24th International Applied Geochemistry Symposium (IAGS 2009) that will be held in Fredericton, New Brunswick (Canada) from June 1st to June 4th, 2009. I am looking forward to getting together with AAG members and others at this meeting next year.

Future articles in upcoming issues of **EXPLORE** will include overviews of the Trinational (Canada-USA-Mexico) soil survey, transport and deposition of colloidal gold, and hydrogeochemical methods for exploration in Western Australia. AAG welcomes Thermo Scientific-Niton as a new Corporate Sponsor of **EXPLORE**. Our **advertisers** are important to **EXPLORE**; please mention seeing their special efforts on our behalf when you talk or correspond with them. For those interested in additional copies of **EXPLORE**, all past issues of **EXPLORE** are available in PDF format on the AAG website. Contributions to **EXPLORE** are welcome anytime, and guidelines for contributors are listed on the inside back page of each issue.

Recently, **EXPLORE** had an enquiry from an AAG member about the cost of mailing **EXPLORE** flat versus folded in half. Folding **EXPLORE** in half would allow it to be mailed in a smaller envelope. The cost savings in postage of the smaller envelope is considerably lessened by the cost that would be incurred to have the newsletter folded. The net savings to **EXPLORE** would only be \$63 per issue by mailing it in a smaller envelope. For now, we will continue to mail out unfolded copies. We appreciate and encourage members to continue to submit ideas or questions to **EXPLORE**.

Beth McClenaghan





Thermo Scientific Niton® XL3 Know your boundaries.

From the frigid Canadian arctic to the brutal heat of the Australian outback, Thermo Scientific Niton XRF Analysers have become the worldwide standard for geochemical field assay in mining exploration.

- Rapid delineation of ore boundaries
- Fast, simple GPS/GIS integration
- High sample throughput and increased sample density over traditional lab methods
- Direct analysis of RAB and RC drilling to dynamically drive exploration programs

Sales & Service Worldwide

Tel. +1 978 670-7460
Fax +1 978 670-7430
E-mail niton@thermofisher.com
www.thermo.com/niton



Real-time geochemical mapping
The Niton Xl3t connects to your
Bluetooth equipped GPS receiver to
store location coordinates with each
reading.



PAGE 4 NUMBER 140 EXPLORE

The geochemistry of acid-saline waters and drainage... continued from page 2

that acidic groundwaters are common throughout the Wheatbelt (Rogers & George 2005; Shand & Degens 2007) has highlighted that acidity and mobilisation of elements, which may be toxic at high concentrations poses a significant risk. A number of projects are being managed by the Cooperative Research Centre for Landscape Environments and Mineral Exploration (CRC LEME) in collaboration with the WA State Government Department of Water, Department of Agriculture and Food and CSIRO Land and Water. The primary aim of these initiatives is to assess the geochemical risks and impacts on receiving environments of deep drainage, and to identify management options for the disposal of acidic waters. This paper outlines the main conclusions of recent investigations (Shand & Degens 2007) including an evaluation of geochemical risks in receiving environments of the Avon catchment into which deep drains discharge.

Occurrence and generation of acid waters in the Wheatbelt

Acid saline groundwaters are present over large areas of the Wheatbelt, but are more common in the east where up to 70% of groundwaters sampled have acidic pH (pH 3 – 4.5 (Fig. 3). Such groundwaters were known to occur




Figure 3. Locations of drain, lake and ground-water sampling sites (classified by sample pH) in the Avon catchment (from Shand & Degens 2007).

prior to the installation of drains showing that drainage is not the main cause of acidification. Acid hypersaline lakes (Fig. 4) are also a relatively common feature of the Wheatbelt, where the lakes represent strongly evaporated groundwater discharge. Acid hypersaline lakes typically occur in low-lying closed basins and in valley bottoms where they form discontinuous chains (Commander *et al.* 2002). Many of the deep drains installed in the Avon catchment of the Wheatbelt to control groundwater levels are acidic,



Figure 4. Acid hypersaline lake in the Wheatbelt near Koorda.

displaying a similar distribution to acid groundwaters (Fig. 3). The significance of increasing discharge of acidic waters for trace metal mobilisation, and the risk that this poses to aquatic ecosystems, should increasing drainage occur, has only recently been investigated (Shand & Degens 2007).

The origin of acidity in the groundwaters is not known in detail, although it is commonly believed that the oxidation of dissolved iron (Fe) plays an important role in generating acidity (Mann 1983; McArthur *et al.* 1991). This reaction may be more important at the margins of Playa Lakes and drains where reduced iron (Fe²⁺), present in Fe-rich groundwater, is oxidised to Fe³⁺, generating acidity (H⁺) in the process:

$$2Fe^{2+} + \frac{1}{2}O_2 + 5H_2O \Leftrightarrow 2Fe(OH)_{3(S)} + 4H^+$$

This model by itself is unlikely to explain the occurrence and preponderance of acid groundwater in many parts of the Wheatbelt, since Fe-rich waters are common in many other non-acidic areas. In addition, acidity is required (consumed) in order to dissolve iron initially by reductive dissolution of oxidised Fe, hence it is difficult to attain the low pH values of

groundwaters without an additional strong source of acidity (e.g. sulfide oxidation as found in acid mine drainage). New models are currently being developed based on fractionating alkalinity, e.g. removing alkalinity by the precipitation of calcite from stored acidity, the latter being stored and transported in the water. Such models may explain e.g. the correlation between calcrete/calcareous soils and gypsum with areas of low pH groundwater.

Geochemistry of the acid waters

Samples were collected from groundwater boreholes, drains and lakes (receiving environments) in the Avon

The geochemistry of acid-saline waters and drainage... continued from page 4

catchment to assess the potential risks that the drains may impose on receiving environments. All samples were filtered in the field through 0.45 μ m membrane filters and either acidified with ultrapure HNO₃ or retained unacidified and stored at 5° C, depending on the analysis performed. Onsite measurements of pH, specific electrical conductance (SEC), temperature and Eh were taken at many sites (when possible). Drain/creek flow was also estimated whenever possible. The concentrations of major and minor cations and sulfate were determined by Inductively Coupled Plasma Optical Emission Spectrometry (ICP-OES), and trace elements by Inductively Coupled Plasma Mass Spectrometry (ICP-MS). Anion analyses were completed using ion chromatography and colorimetry (APHA 1998).

Drain flow was highly variable (from 30 l s⁻¹ to less than 0.1 l s⁻¹) over the study area, and generally decreased with increasing age of the drain. In some cases, drains ceased to flow once the water table fell below the base of the drains. However, the impact often extended to distances greater than 200 m (Ali et al. 2004a) depending on the local geology and hydrogeology. During periods of low flow, the salinity of drain water increased and pH generally decreased.

The pH data for surface waters and groundwater

displays a bimodal distribution (Fig. 5). The pH was generally lower in the eastern Wheatbelt than in the western and central Wheatbelt, and most drains were typically saline (Fig. 5). Major element concentrations and TDS display large variations, typically over several orders of magnitude. High TDS surface waters are present in both acidic and alkaline surface waters and groundwaters (Fig. 6), although the former have much higher TDS, largely due to evaporation. Sodium displays a strong positive correlation with Cl (r^2 =0.99) but the other major elements correlate poorly with Cl indicating gains or losses relative to seawater (Fig. 7). The surface waters in particular have low ratios of Ca, Mg and SO₄ to Cl relative to seawater consistent with removal processes such as mineral precipitation (e.g. calcite, gypsum, halite) and/or removal through redox processes (e.g., reduction of sulfate to sulfide).

Most minor and trace elements vary over several orders of magnitude. Nearly 50% of the samples have F concentrations exceeding the WHO guideline value of 1.5 mg l⁻¹ (Fig. 8). Nutrients and total organic carbon were measured only in surface waters but reached high concentrations: PO₄-P, NO₃-N, NH₄-N and TOC had maxima of 1.4, 11, 45

continued on page 6




Figure 5. Boxplots (displaying median, interquartile ranges and outliers) and histograms for TDS (left) and pH (right) in surface waters and groundwaters sampled in the Avon catchment.

PAGE 6 NUMBER 140 EXPLORE

The geochemistry of acid-saline waters and drainage... continued from page 5




Figure 6. Relationship of TDS and major element concentrations to pH in surface waters and groundwaters of the Avon catchment.




Figure 7. Relationship of TDS, pH and major element concentrations to Cl in surface waters and groundwaters of the Avon catchment. Solid line is seawater dilution line.

and 255 mg l⁻¹, respectively. The redox-sensitive elements Fe and Mn are present at high concentration in the waters, in general being higher at lower pH. Aluminium concentrations also display a wide range, reaching maximum concentrations greater than 1000 mg l⁻¹ at low




Figure 8. Concentrations of minor and trace elements in waters of the Avon catchment plotted against pH. Note log scales for Fe, Mn and Al.

pH. Many metals, including Fe and Mn are more soluble and mobile under acidic conditions (Stumm & Morgan 1996). The acid nature of both surface waters and groundwaters has allowed a number of trace elements to reach mg l⁻¹ concentrations including Co, Cr, Cu, Ni, Pb, U, Y and the rare earth elements (REE), many above national drinking water guideline values (Fig. 8). A number of elements which are rarely present in waters such as Th, Pd and Tl have elevated concentrations above typical baseline values. The concentrations of many trace elements are more typical of highly polluted waters, and much higher than those expected in non-industrial or mining areas (Nordstrom 2007; Shand & Edmunds 2008).

Geochemical processes occurring in drains

The discharge of acid groundwater to artificial drains, and subsequent reactions, has led to a complex and dynamic geochemical environment. The oxidation of reduced soluble Fe²⁺ in drains to insoluble Fe³⁺ oxy-hydroxides is commonly strikingly visible (Fig. 9) and often lead to a



Figure 9. Iron oxy-hydroxide minerals covering the base of a stream, note also the abundance of salt efflorescences adjacent to the stream caused by evaporation of capillary water. The efflorescences are generally present during summer months but are washed into the stream during rainfall periods, producing a pulse of increased salinity.

The geochemistry of acid-saline waters and drainage... continued from page 6

further decrease in pH. The elements Fe and Al are pH sensitive and act as stores of acidity in solution, releasing H⁺ during the precipitation of oxyhydroxide minerals by hydrolysis reactions. These minerals commonly form gels in the environment, which are efficient scavengers of trace metals (e.g., As, Cr, Sn, Th, V, Cd, Cu, Mo), but the fine colloidal nature of these materials places them at risk of being suspended and flushed from the drains even at low flow velocities. The association of the metals with the precipitates and gels is not well documented in the drains of the study area, but it is probable that these materials could act as transporting media for metals from the drains if not well managed.

A wide range of minerals actively form in the drain environment (Fitzpatrick et al. 2007). Iron sulfide minerals (pyrite and monosulfides) are commonly encountered immediately beneath the oxidised sediment surface where conditions are reducing, causing sulfate to be reduced to sulfide. The key requirements for high rates of sulfate reduction and sulfide accumulation are: (i) high concentrations of sulfate in surface water or groundwater, (ii) saturated, reducing iron-rich soils and sediments, and (iii) the availability of labile carbon to fuel microbial activity (Berner 1984). Disturbance of these sulfide-rich sediments, in particular if they form monosulfidic black oozes, in drains can lead to severe deoxygenation and acidification. (Sullivan et al. 2002).

A remarkable range of other mineral precipitates and gels were found in the drains (Table 1), indicative of the

thenardite NaSO ₄	starkeyite MgSO ₄ ·4(H ₂ O)	pentahydrite MgSO ₄ ·5(l rozenite Fe ²⁺ SO ₄ ·4	Mineral Name Chemical Formula	akaganéite bassanite barite bischofite bloedite boehmite carnallite eugsterite glauberite gypsum halite jarosite lepidocrocite mirabilite natrojarosite pentahydrite rozenite schwertmannite starkeyite	Fe ³⁺ O(OH, CI) 2(Ca) ₂ (SO ₄)·(H ₂ O) BaSO ₄ MgCl ₂ ·6H ₂ O Na ₂ Mg(SO ₄) ₂ ·4H ₂ O AlO(OH) KMgCl ₃ ·6(H ₂ O) Na ₄ Ca(SO ₄) ₃ ·2(H ₂ O Na ₂ Ca(SO ₄) ₂ CaSO ₄ ·2(H ₂ O) NaCl KFe ³⁺ (SO ₄) ₂ (OH) ₆ FeO(OH) Na ₂ SO ₄ ·10(H ₂ O) Na Fe ³⁺ (SO ₄) ₂ (OH) MgSO ₄ ·5(H ₂ O) Fe ²⁺ SO ₄ ·4(H ₂ O) Fe ³⁺ ₁₆ O ₁₆ (OH) ₁₂ (SO MgSO ₄ ·4(H ₂ O)
pentahydrite $MgSO_4 \cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4 \cdot 4(H_2O)$ schwertmannite $Fe^{3+}_{16}O_{16}(OH)_{12}(SO)$	pentahydrite $MgSO_4 \cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4 \cdot 4(H_2O)$		bassanite barite barite bischofite bloedite boehmite carnallite glauberite gypsum halite jarosite basSO ₄ MgCl ₂ ·6H ₂ O MgCl ₂ ·6H ₂ O MgCl ₃ ·4H ₂ MgCl ₃ ·6(H ₂ O) MgCl ₃ ·6(H ₂ O) Na ₂ Mg(SO ₄) ₂ ·4H ₂ Na ₂ Ca(SO ₄) ₃ ·2(H ₂ O) Na ₄ Ca(SO ₄) ₃ ·2(H ₂ O) NaCl KFe ³⁺ (SO ₄) ₂ (OH)	mirabilite	Na ₂ SO ₄ ·10(H ₂ O)
mirabilite $Na_2SO_4 \cdot 10(H_2O)$ natrojarosite $Na Fe^{3+}(SO_4)_2(OH)$ pentahydrite $MgSO_4 \cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4 \cdot 4(H_2O)$ schwertmannite $Fe^{3+}_{16}O_{16}(OH)_{12}(SO)$	mirabilite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na Fe^{3+}(SO_4)_2(OH)$ pentahydrite $MgSO_4\cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4\cdot 4(H_2O)$		bassanite $2(Ca)_2(SO_4) \cdot (H_2O)$ barite $BaSO_4$ bischofite $MgCl_2 \cdot 6H_2O$ bloedite $Na_2Mg(SO_4)_2 \cdot 4H_2O$ boehmite $AIO(OH)$ carnallite $KMgCl_3 \cdot 6(H_2O)$ eugsterite $Na_4Ca(SO_4)_3 \cdot 2(H_2O)$ glauberite $Na_2Ca(SO_4)_2$ gypsum $CaSO_4 \cdot 2(H_2O)$	jarosite	KFe ³⁺ (SO ₄) ₂ (OH) ₆
jarosite $KFe^{3+}(SO_4)_2(OH)_6$ lepidocrocite $FeO(OH)$ mirabilite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na Fe^{3+}(SO_4)_2(OH)$ pentahydrite $MgSO_4\cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4\cdot 4(H_2O)$ schwertmannite $Fe^{3+}_{16}O_{16}(OH)_{12}(SO)$	jarosite $KFe^{3+}(SO_4)_2(OH)_6$ lepidocrocite $FeO(OH)$ mirabilite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na_2SO_4\cdot 10(H_2O)$ pentahydrite $MgSO_4\cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4\cdot 4(H_2O)$	jarosite KFe³+(SO₄ lepidocrocite FeO(OH)	bassanite $2(Ca)_2(SO_4) \cdot (H_2O)$ barite $BaSO_4$ bischofite $MgCl_2 \cdot 6H_2O$ bloedite $Na_2Mg(SO_4)_2 \cdot 4H_2O$ boehmite $AIO(OH)$ carnallite $KMgCl_3 \cdot 6(H_2O)$ eugsterite $Na_4Ca(SO_4)_3 \cdot 2(H_2O)$	gypsum	CaSO₄·2(H₂O)
gypsum $CaSO_4 \cdot 2(H_2O)$ halite $NaCl$ jarosite $KFe^{3+}(SO_4)_2(OH)_6$ lepidocrocite $FeO(OH)$ mirabilite $Na_2SO_4 \cdot 10(H_2O)$ natrojarosite $Na_2SO_4 \cdot 10(H_2O)$ pentahydrite $MgSO_4 \cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4 \cdot 4(H_2O)$ schwertmannite $Fe^{3+}_{16}O_{16}(OH)_{12}(SO_4)$	gypsum halite halite jarosite lepidocrocite mirabilite natrojarosite pentahydrite rozenite SaSO ₄ ·2(H ₂ O) NaCl KFe³+(SO ₄) ₂ (OH) ₆ FeO(OH) Na ₂ SO ₄ ·10(H ₂ O) Na Fe³+(SO ₄) ₂ (OH) MgSO ₄ ·5(H ₂ O) Fe²+SO ₄ ·4(H ₂ O)	gypsum CaSO₄·2(h halite NaCl jarosite KFe³+(SO₄ lepidocrocite FeO(OH)	bassanite $2(Ca)_2(SO_4) \cdot (H_2O)$ barite $BaSO_4$ bischofite $MgCl_2 \cdot 6H_2O$ bloedite $Na_2Mg(SO_4)_2 \cdot 4H_2O$ boehmite $AIO(OH)$	· ·	
glauberite $Na_2Ca(SO_4)_2$ gypsum $CaSO_4 \cdot 2(H_2O)$ halite $NaCl$ jarosite $KFe^{3+}(SO_4)_2(OH)_6$ lepidocrocite $FeO(OH)$ mirabilite $Na_2SO_4 \cdot 10(H_2O)$ natrojarosite $Na Fe^{3+}(SO_4)_2(OH)_6$ pentahydrite $MgSO_4 \cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4 \cdot 4(H_2O)$ schwertmannite $Fe^{3+}_{16}O_{16}(OH)_{12}(SO_4)$	$\begin{array}{cccccccccccccccccccccccccccccccccccc$	glauberite Na₂Ca(SO gypsum CaSO₄·2(H halite NaCl jarosite KFe³+(SO₄ lepidocrocite FeO(OH)	bassanite $2(Ca)_2(SO_4) \cdot (H_2O)$ barite $BaSO_4$ bischofite $MgCl_2 \cdot 6H_2O$ bloedite $Na_2Mg(SO_4)_2 \cdot 4H_2O$		\ /
carnallite KMgCl ₃ ·6(H ₂ O) eugsterite Na ₄ Ca(SO ₄) ₃ ·2(H ₂ glauberite Na ₂ Ca(SO ₄) ₂ gypsum CaSO ₄ ·2(H ₂ O) halite NaCl jarosite KFe ³⁺ (SO ₄) ₂ (OH) ₆ lepidocrocite FeO(OH) mirabilite Na ₂ SO ₄ ·10(H ₂ O) natrojarosite Na Fe ³⁺ (SO ₄) ₂ (OH pentahydrite MgSO ₄ ·5(H ₂ O) rozenite Fe ²⁺ SO ₄ ·4(H ₂ O) schwertmannite Fe ³⁺ 1 ₆ O ₁₆ (OH) ₁₂ (SO	carnallite $KMgCI_3 \cdot 6(H_2O)$ eugsterite $Na_4Ca(SO_4)_3 \cdot 2(H_2)$ glauberite $Na_2Ca(SO_4)_2$ gypsum $CaSO_4 \cdot 2(H_2O)$ halite $NaCI$ jarosite $KFe^{3+}(SO_4)_2(OH)_6$ lepidocrocite $FeO(OH)$ mirabilite $Na_2SO_4 \cdot 10(H_2O)$ natrojarosite $Na Fe^{3+}(SO_4)_2(OH)_6$ pentahydrite $MgSO_4 \cdot 5(H_2O)$ rozenite $Fe^{2+}SO_4 \cdot 4(H_2O)$	carnallite KMgCl ₃ ·6(eugsterite Na ₄ Ca(SO glauberite Na ₂ Ca(SO gypsum CaSO ₄ ·2(H halite NaCl jarosite KFe ³⁺ (SO ₄ lepidocrocite FeO(OH)	bassanite $2(Ca)_2(SO_4) \cdot (H_2O)$		Na ₂ Mg(SO ₄) ₂ ·4H ₂ 0
bloedite $Na_2Mg(SO_4)_2\cdot 4H_2O_5$ boehmite $AIO(OH)$ carnallite $KMgCl_3\cdot 6(H_2O)$ eugsterite $Na_4Ca(SO_4)_3\cdot 2(H_2O_5)$ gypsum $CaSO_4\cdot 2(H_2O)$ halite $NaCl$ jarosite $KFe^{3+}(SO_4)_2(OH)_6$ lepidocrocite $FeO(OH)$ mirabilite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na_2SO_4\cdot 10(H_2O)$ rozenite $Na_2SO_4\cdot 10(H_2O)$	bloedite $Na_2Mg(SO_4)_2\cdot 4H_2O_4$ boehmite $AIO(OH)$ carnallite $KMgCl_3\cdot 6(H_2O)$ eugsterite $Na_4Ca(SO_4)_3\cdot 2(H_2O_4)_2$ glauberite $Na_2Ca(SO_4)_2$ gypsum $CaSO_4\cdot 2(H_2O)$ halite $NaCl$ jarosite $KFe^{3+}(SO_4)_2(OH)_6$ lepidocrocite $FeO(OH)$ mirabilite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na_2SO_4\cdot 10(H_2O)$ natrojarosite $Na_2SO_4\cdot 10(H_2O)$ pentahydrite $Na_2SO_4\cdot 10(H_2O)$	bloedite boehmite carnallite eugsterite glauberite gypsum halite jarosite lepidocrocite Na2Mg(SC AIO(OH) NA4Ca(SO NA4Ca(SO RA4Ca(SO RA50) NA2Ca(SO RA50) NA2Ca(SO RA50) NA2Ca(SO RA50) NACI SA50 NA		akaganéite bassanite	Fe³+O(OH, CI) 2(Ca) ₂ (SO ₄)·(H ₂ O)

Table 1 Chemical compositions of minerals found in the drains of the Australian Wheatbelt.

continued on page 8



PAGE 8 NUMBER 140 EXPLORE

The geochemistry of acid-saline waters and drainage... continued from page 7

range of geochemical conditions occurring. Bright yellow natrojarosite mottles in some of the clay-rich sulfuric horizons are indicative of acid conditions in the pH range 3.5-4. Similarly, the occurrence of orange-coloured mottles, gels and crusts are indicative of schwertmannite and akaganéite, which form from the oxidation of ferrous iron under acidic conditions in the pH range 4-5. Saline and subaqueous soils with sulfuric material may also occur in receiving lakes. Other precipitates include evaporite minerals, those being found to date in the drains and lakes include pentahydrite, starkeyite, bischofite, bassanite, carnallite, rozenite, barite, halite and gypsum in sandy sulfuric horizons with pH <3.0; natrojarosite and jarosite in clay-rich sulfuric horizons with pH 3.5-4; and eugsterite, bloedite, thenardite, glauberite, gypsum, thenardite, mirabilite, schwertmannite, lepidocrocite, akaganéite and colloidal poorly crystalline, pseudoboehmite-like (white) precipitates in sulfidic materials with pH >4-5.

The processes that give rise to the variable mineral assemblages play a major role in influencing transport of acidity and trace metals from the drains and vary with season and drain age. Furthermore, identifying the processes has enabled development of practical management options to mitigate the risks that drain discharge poses to receiving environments. In order to facilitate the transfer of information on drain and water geochemistry, an interactive web-based database has been constructed (Baker & Fitzpatrick 2005). This website incorporates a visual









CANADA
MEXICO
CHILE
BRAZIL
ARGENTINA
ECUADOR
PERU
GUYANA
VENEZUELA
TURKEY



New Vancouver Shipping Address

1020 Cordova Street East, Vancouver, BC Canada V6A 4A3 Phone 604 253 3158 Fax 604 253 1716 Email acmeinfo@acmelab.com

WWW.ACMELAB.COM

aid designed to allow rapid and effective communication of data and information at a range of scales (regional to microscopic).

Receiving environments

Many acid drains discharge to saline sites, principally playas in the floodways of the main palaeodrainage systems (Dogramaci & Degens 2003; de Broekert & Coles 2004; Ali et al. 2004b). Acid drainage discharge to playa lakes can result in acidification of surface waters and the sediments and soils surrounding the playa lakes. However, in some cases lake waters and sediments may have been acidic prior to discharge, in which case the impacts are dependent on the magnitude of drain discharge compared with regional ground-water discharge. Concentrations of Al, Fe and trace metals including Pb, Ni, Co and U are often higher in the surface waters of sites receiving acidic drainage. This acidification and associated increased in trace metal solubility may result in impacts on aquatic ecosystems, including loss of habitat and reduced ecosystem functioning during playa lake filling events, and should form an integral part of future research in the region.

The discharge of acid drain water into alkaline waters in receiving environments allows the possibility for neutralisation of acidity. Geochemical modelling using the PHREEQC geochemical code was undertaken. This modelling indicates that the neutralisation of acidity in the drain waters in receiving environments is unlikely to occur until the volumes of alkaline waters are in the order of more than 25-99 times that of the incoming acid drainage waters. It will therefore be necessary to implement further remediation strategies in the absence of such volumes of alkaline waters being continuously available.

Management implications of acid drain water

The widespread occurrence of acid groundwaters and surface waters in the Wheatbelt containing high concentrations of potentially toxic elements requires care to be taken in managing and linking deep drains installed for salinity reduction. Drains also need to be designed and managed to minimise turbulent flow to minimise the flushing of precipitates and gels (frequently containing trace metals) and disturbance of sulfidic sediments (being a store of acidity and trace metals). In particular, entry of surface waters from catchments to the drains should be avoided without measures to contain flow velocities. Drains should also be designed to maximise hydrological residence times as the formation of precipitates will contribute to maximising retention of trace metals within the systems.


Management of trace metal mobility and acid release will need to be considered when maintenance cleaning of sediments from drains is carried out. This might include mixing with sediments from alkaline drain spoils, placement within depressions on drain spoils (allowing drying and containment but including contact with alkaline spoils) or collection and containment in a site with low risk of off-site impacts (i.e. outside of a surface water flow path).

ALS Laboratory Group ANALYTICAL CHEMISTRY & TESTING SERVICES

Mineral Division - ALS Chemex











e are a global analytical testing company serving the mining and mineral exploration industry in 17 countries around the world.

We specialise in:

- Assaying and geochemical analysis for mining and mineral exploration projects
- On-site contract mine laboratory management
- · Environmental monitoring for mining operations
- · Acid mine drainage studies

Contact the ALS Chemex laboratory nearest to you to find out how we can help you with all of your project's analytical needs. Detailed contact information for individual locations can be found on our website.

www.alschemex.com

Right solutions....
Right partner

NUMBER 140 EXPLORE PAGE 10

The geochemistry of acid-saline waters and drainage... continued from page 8

It is vital for all landholders, community groups, drainage contractors and local governments to be aware of the many impacts that could result from the occurrence of sulfuric materials and disturbance of sulfidic materials, as these have important consequences for environmental, engineering, economic, and quality of life perspectives. Disturbance and oxidation of sulfidic material can have negative impacts including: destruction of wetlands, acidification and deoxygenation of waterways and increased incidence of fish kills and disease, contamination of valuable groundwater resources, mobilisation and accumulation of heavy metals, corrosion and destabilisation of roads, concrete and steel infrastructure, stimulation of blooms of blue-green algae, decrease in the agricultural productivity of land, increased odour problems and increased mosquito and arbovirus incidence.

Future work

The work completed in the Avon catchment is now being extended to other parts of the Wheatbelt. Following a review of potential acid neutralisation techniques for acid drains in the Wheatbelt (Douglas & Degens 2006), pilot treatment sites have recently been established to investigate low-cost, practical treatment options for acidic drainage waters. Future research will focus on the geochemical processes and efficiency of the treatment sites. The trials will focus on simple engineering structures using either carbonate neutralisation (being the most widely available, low cost

material currently available in the Wheatbelt) or sulfate reduction systems using locally available carbon sources.

Acknowledgements

We are grateful to the following team members of the project for their scientific contributions: Andrew Baker, Grant Douglas, Rob Fitzpatrick, Richard George, David Gray, Warren Hicks, Adam Lillicrap, Mark Raven and Margaret Smith.

References

- Ali, R., Hatton, T., George, R., Byrne, J. & Hodgson, G., 2004a. Evaluation of the impacts of deep open drains on groundwater levels in the Wheatbelt of Western Australia. Australian Journal of Agricultural Research, 55, 1159-1171.
- Ali, R., Hatton, T., Lambert, T., Byrne, J., Hodgson, G. & George, R. 2004b. An assessment of the quality and quantity of discharge from deep open drains in Narembeen: Wheatbelt of Western Australia. In: 1st National Salinity Engineering Conf., 9-12 Nov., 2004, Perth Australia, 84-88.
- Allison, G. B., Cook, P. G., Barnett, S. R., Walker, G. R., Jolly, I. D & Hughes, M. W. 1990. Land clearance and river salinisation in the Western Murray Basin. Journal of Hydrology, 119, 1-20
- APHA, 1998. Methods for the examination of water and waste water, 20th edition, published jointly in 1998 by the American Public Health Association, American Water Works Association and the Water Environment Federation. Method 3120 for cations, method 4110 anions. continued on page 11

Intertek Minerals Services

Offering Analytical Laboratory & Sample Preparation Services

- Australia •Indonesia South Africa
- •China PhilippinesPNG

www.intertek-cb.com/minerals







Robotic Laboratories



The geochemistry of acid-saline waters and drainage... continued from page 10

- Baker, A.K.M. & Fitzpatrick R.W. 2005. A systematic web-based approach for the acquisition, collation and communication of soil-regolith data. In: Roach, I.C. (ed.) Regolith 2005 - Ten Years of CRC LEME. CRC LEME, Perth, Australia, 3-7.
- Berner, R.A. 1984. Sedimentary pyrite formation: an update. Geochimica et Cosmochimica Acta, 48, 605–615.
- Clarke, C.J., George, R.J., Bell, R.W. & Hatton, T.J. 2002. Dryland salinity in South-Western Australia: its origins, remedies and future research directions. Australian Journal of Soil Research, 40, 93-113.
- Commander P., Schnoknecht N., Verboom B. & Cacetta P. 2002. The geology, physiography and soils of Wheatbelt valleys. In: Read, V. & Associates (eds), Dealing with salinity in Wheatbelt valleys: Processes, prospects, and practical options, Water and Rivers Commission, Perth. 1-27.
- De Broekert, P.P. & Coles, N.A. 2004. Wetland basins for saline drainage water disposal: Bodallin and Elachbutting Catchments, Eastern Wheatbelt, Western Australia. Department of Agriculture WA, Resource Management Technical Report 290. October 2004.
- Dogramaci S. & Degens B. 2003. Review of Engineering and Safe Disposal Options, Western Australia. Water and Rivers, Report No. SLUI 20.
- Douglas, G. & Degens, B. 2006. A synopsis of potential amendments and remediation techniques for the neutralization of acidic drainage waters in the Western Australian Wheatbelt. CRC LEME Open File Report 209, 22
- Fitzpatrick, R., Degens, B., Baker, A., Raven, M., Smith, M., Rogers, S., George R. & Shand, P. 2007. Geochemical and hydrochemical processes occurring during drainage of saline watertables in the Avon Basin. In: P. Shand, P. & B. Degens (eds.) Avon Catchment Acid Groundwater - Geochemical Risk Assessment. CRC LEME Open File Report 191.
- Hatton, T.J., Ruprecht, J. & George, R.J. 2003. Preclearing hydrology of the Western Australia Wheatbelt: Target for the future? Plant and Soil, 257, 341-356.
- Mann, A.W. 1983. Hydrogeochemistry and weathering on the Yilgarn Block, Western Australia – ferrolysis and heavy metals in continental brines. Geochimica et Cosmochimica Acta, 47, 181-190.
- McArthur, J.M., Turner, J.V., Lyons. W.B., Osborn, A.O. & Thirlwall, M.F. 1991. Hydrochemistry on the Yilgarn Block, Western Australia: Ferrolysis and mineralization in acidic brines. Geochimica et Cosmochimica Acta, 55, 1273-1288.
- Nordstrom, D.K. 2007. Questa baseline and pre-mining groundwater quality investigation. 25. Summary of results and baseline and pre-mining ground-water geochemistry, Red River Valley, Taos County, New Mexico, 2001-2005: U.S. Geological Survey Professional Paper 1728, 170 pp.
- Rogers, S.L. & George, R. 2005. WA Wheatbelt drainage acidic groundwater, not just a salt issue, Focus on Salt, Issue 33, June 2005, 8-9.
- Shand, P. & Degens, B. (eds.) 2007. Avon Catchment Acid Groundwater - Geochemical Risk Assessment. CRC LEME Open File Report 191.

- Shand, P. & Edmunds, W.M. 2008. Baseline Inorganic Quality of Groundwaters. In: W.M. Edmunds & P. Shand (eds.), Natural Groundwater Quality. Blackwell, Oxford.
- Stumm, W. & Morgan, J.J. 1996. Aquatic Chemistry. Wiley & Sons, New York.
- Sullivan, L.A., Bush R.T. & Fyfe, D. 2002. Acid sulfate soil drain ooze: distribution, behaviour and implications for acidification and deoxygenation of waterways. In: Lin, C., Melville, M.D. & Sullivan, L.A. (eds.) Acid sulfate soils in Australia and China. Science Press, Beijing. 91-99.

Paul Shand

CSIRO Land and Water/CRC LEME, Private Bag 2, Glen Osmond, SA 5064, Australia.

E-mail: Paul.shand@csiro.au

Brad Degens

Water Resource Management Division, Department of Water WA, P.O. Box K822 Perth, WA 6842, Australia. E-mail: Brad.degens@water.wa.gov.au

Steve Rogers

CRC LEME, CSIRO Exploration and Mining, 26 Dick Perry Avenue, Kensington WA 6151, Australia E-mail: Steve.rogers@csiro.au

A rock solid reputation for absolute accuracy

ince 1982, Becquerel Laboratories has been meeting the analytical requirements of clients in the geological and mineral exploration sector.

Our company is a world leader in Neutron Activation Analysis (NAA). An ultra-sensitive technique, NAA is an ideal quality control procedure to complement and verify results from other analytical techniques. It provides cost-effective, timely analysis and one simple method can analyze over 30 separate elements.

For absolute accuracy, trust the analytical expertise of Becquerel Laboratories.

For more information call 905-826-3080 or visit www.becquerellabs.com



Accredited to ISO 17025 by the Standards Council



6790 Kitimat Road, Unit 4 Mississauga, Ontario Canada L5N 5L9

absolute accuracy

PAGE 12 NUMBER 140 EXPLORE

The United Nations International Year of Planet Earth (2008)

2008 INTERNATIONAL YEAR OF PLANET EARTH Earth Sciences for Society

The United Nations International Year of Planet Earth (2008) global launch was held at UNESCO Headquarters (Paris, France) on February 12 and 13, 2008 (Fig. 1). The



Figure 1. UNESCO flag waving beside the UN headquarters "World Dome" with the Eiffel Tower in the background.

International Year of Planet Earth (IYPE) is designed to foster scientific research activities and outreach, by raising global political and public awareness of the immense, under-utilized potential of Earth sciences for improving the quality of life and safeguarding the planet. International Union of Geological Sciences (IUGS) and UNESCO are the sponsoring partners of IYPE, with all 191 UN member countries supporting the UN Resolution proclaiming the IYPE for 2008. The resolution states "The International Year of Planet Earth aims to capture people's imagination with the exciting knowledge we possess about our planet, and to see that knowledge used to make the Earth a safer, healthier and wealthier place for our children and grandchildren International Year of Planet Earth 2007-2009."

The founding patrons of IYPE are the President of Namibia, Sam Nujoma; the former President of Tanzania,





Benjamin Mkapa; the Chairman of the Board of Anglo American, Sir Mark Moody-Stuart; the former Prime Minister of the Netherlands, Ruud Lubbers; and King Carl XVI Gustaf of Sweden. By the official launch, there were approximately 70 member nations, including Canada. The ceremonies were dominantly a series of incredible speeches and theme lectures (Fig. 2), and amazing poems from young people around the world. These presentations were eclipsed by a speech by Sir Arthur C. Clarke that was definitely the highlight of the IYPE launch ceremonies. Clarke stated: "The aim of the Year is to persuade public and governments worldwide to make better use of Earth science when framing planning decisions, and by using Earth science to inform the



Figure 2. USGS Director Mark Myers presenting a talk (with a discussion panel) about the "evolution of humans" with the backdrop of a NASA composite photographic image of Earth's night sky (lit up), by far achieved by fossil fuel comsumption and not considered sustainable.

public about the sustainable use of Earth resources, to make the world a healthier, wealthier and safer place in which to live."

Almost 70 countries are members of IYPE. The two main activities of IYPE are its Science and Outreach programs. Funding for projects in these 2 areas is to come from industry, foundations and governments around the world, although implementing them has mainly been at the grassroots level by volunteers With greater than 400,000 earth scientists worldwide, the UN is hoping we have a broad-based impact. The Science Programs fall into 10 broad, societal relevant and multidisciplinary themes:

continued on page 13

The United Nations International Year of Planet Earth... continued from page 12

health, climate, groundwater, ocean, soils, deep Earth, megacities, hazards, resources, and life. Efforts also include an IYPE book on-line that has lots of "gift ideas". http://yearofplanetearth.org

In January 2008, I asked the AAG executive for input before I traveled as the AAG representative to the IYPE launch (Fig. 3) Here is a summary of some of the feedback I received. AAG members are actively engaged in outreach in schools, as are many other active geoscience organizations. Because AAG is an international organization, however, the impacts of the various Science components of IYPE were of most interest to us asour expertise falls in over half of the Science theme areas. Our AAG members plan on



Figure 3. Dave Lentz, the AAG representative at the United Nations International Year of Planet Earth (2008) global launch at UNESCO Headquarters (Paris, France) on February 12 and 13, 2008.

being engaged in them all from grassroots efforts through to lobbying industry and governments, particularly on issues pertaining to world health and poverty. These efforts can be achieved by encouraging the broad-based application of environmental and exploration geochemistry. As major proponents of detailed regional geochemical mapping, all AAG members know we can help to identify areas where potentially toxic elements can negatively impact health, where one can see the effect of industrial or natural pollution and perhaps link it to health problems, particularly cancers and malnutrition; however, this occurs only where geochemical data are available for epidemiological studies. Numerous cases exist where surveys have led to awareness of health problems related to a wide variety of heavy metals. The FOREGS Geochemical Baseline Mapping Program for sediment, water, and soil recognized and addresses these issues by doing the research, building the protocols, and then building the databases covering much of Europe. IUGS-IAGC's Working Group on Global Geochemical Baselines is also contributing to this important effort; there are numerous websites that detail these baseline studies and method development, and that determine protocols for sampling through to analysis. In 2006, the North American Soil Geochemical Landscapes Project was initiated and represents a comprehensive tri-national soil sampling effort involving multiple agencies; they have a special session at the IAGS 2009.

As we know, these same regional geochemical databases and maps catalyze mineral exploration and, in rare cases, may result in mine development, which can have a major impact on poverty. Recently published statistics from northern Ontario (Canada) quantify the huge economic impact mining has had in generating wealth. Having just spent a few weeks in Mali at a gold mine, I saw that this impact was overtly evident; this year alone yielded profits to the government of approximately 80 million dollars. Mali is considered to be one of the five poorest countries in the world, with >60% of its population earning less than \$1 USD per day. Much of the Mali's government funds are spent on health and clean water development.

Dr. Olle Selinus (Geological Survey of Sweden) has long been working on issues related to world health and poverty, in Sweden and around the Third World. For some time, Ollie's group has been focused on medical geology in developing nations. The International Medical Geology Association (IMGA) is focused on these issues; many AAG members are actively engaged in this field too, where the lines between disciplines fade to white. Presently, the IMGA is developing a web-based education package for general use.

Public awareness will lead to government attention, which is what is needed. In Canada for example, the Canadian International Development Agency (CIDA) does not generally support these types of programs, even though the long term impacts are known to be high. CIDA's focus in Mali is on clean drinking water and health too, but to many they do not see the commonality of these strategies.

AAG has a very engaged membership. There is a lot of volunteer work ahead of us to make the level of impact that only we can bring to the world. A major grassroots effort focused on public understanding through to motivating government leadership and participation is needed, as we have seen with the British Geological Survey and Swedish Geological Survey. We really can change the world, if we really want to.

More information about the United Nations International Year of Planet Earth (2008) is available on the following website: http://yearoftheplanetearth.org.

David Lentz

Department of Geology, University of New Brunswick 2 Bailey Drive, Box 4400, Fredericton, NB E3B 5A3 CANADA Email: dlentz@unb.ca



PAGE 14 NUMBER 140 EXPLORE



24TH INTERNATIONAL APPLIED GEOCHEMISTRY SYMPOSIUM (IAGS 2009)


WELCOME

The 24th International Applied Geochemistry Symposium (IAGS 2009) is the Association of Applied Geochemists (AAG) biennial meeting, which will be held in Fredericton (New Brunswick, Canada) on the campus of the University of New Brunswick Monday June 1st to Thursday June 4th, **2009.** This biennial **AAG** meeting is co-sponsored by the International Association of GeoChemistry (IAGC) and the International Association of GeoAnalysts (IAG) and includes the North Atlantic Minerals Symposium (NAMS). The symposium will be preceded by 5 professional development workshops on Sunday, May 31st. There will be 3 premeeting field trips (Wednesday May 27 to Saturday May 30th) and 3 post-meeting field trips (Friday June 5th to Monday June 8th) throughout Atlantic Canada. The trips will leave from and return to Fredericton. The symposium is jointly organized by geoscientists from the *University* of New Brunswick (UNB), New Brunswick Department of Natural Resources, New Brunswick Research and Productivity Council, New Brunswick Department of the Environment, and professionals drawn from the consulting engineering and mineral exploration industry in New Brunswick, in conjunction with a professional conference organizer (PCO).

TECHNICAL PROGRAM

The technical program of the symposium will run 4 days and will include 2 separate morning ½ day Plenary Sessions (1 Mineral Exploration oriented and the 2nd Environmentally oriented) that will feature a keynote speaker and will not overlap with other sessions. The remaining days will have ½ day sessions with 15-minute presentations. Tuesday afternoon will feature 5-minute oral presentations (4 slide) for the poster presenters.

An extended abstract volume will be produced with a format for each paper of 2-4 pages, a 200 word short abstract, figures, tables, acknowledgements, and references in a Microsoft Word document. An example of the submission format will be available on the IAGS 2009 website soon.



These extended abstracts will be printed in a volume and reproduced on CD-ROM; student registrants will only receive the CD-ROM. Both the book and CD-ROM will be included in the normal registration package. The co-chairs will be involved with the scientific and technical editing of these extended abstracts, with the help of the Technical Program Chairs, Drs. Kay Thorne (NB DNR-Minerals) and David Keighley (UNB).

Abstracts will be accepted starting January 15th, 2009 Submission will be by email to: dlentz@unb.ca. **The final deadline for extended abstracts is February 15th, 2009**

PLENARY SESSION IN HONOUR OF PROFESSOR GERRY GOVETT

Deep Search Geochemical Exploration Methods
Chair: Wayne Goodfellow (Geological Survey of Canada)
Dr. Gerry Govett is one of the pioneers in the development and testing of new and improved methods of detecting ore deposits buried at depth in the Earth's crust. As leader of the Exploration Geochemistry Group at the University of New Brunswick (UNB), Professor Govett led a team of researchers and graduate students focused on primary halos associated with massive sulphide deposits, the electrochemical dispersion of elements in soils overlying buried deposits, and the mathematical processing of geochemical data to discern trends related to mineralized sources.

With the return of the 24th International Applied Geochemical Symposium (IAGS) to Canada in 2009, a plenary one-day session has been planned to recognize Dr. Govett's early contributions to the development of deep search geochemical methods. The focus on deeply penetrating methods recognizes the increasing need to replenish the declining base metal reserves of major mining camps in Canada and indeed elsewhere in the world, and the need for more effective geochemical methods of detecting deposits concealed beneath thick glacial sediment or at depth within rock sequences. The session will consist of talks and posters on a range of topics related to deep exploration. These include primary mineralogical and geochemical vectors to ore such as hydrothermal alteration and dispersion of elements and isotopes from vents in the case of seafloor hydrothermal deposits; secondary vectors related to mineral and element dispersion by a number of processes such as glaciation, ground and surface water transport and deposition of metals and tracer isotopes; the upward migration of elements into soils by advective or electrochemical mechanisms; novel methods employing breakthroughs in analytical technology; and 3D GIS visualization and interpretation of prospective ore horizons and related primary vectors. The objective of this session is to present recent advancements in deep search methods. The oral program will consist of both invited speakers to cover the major themes and unsolicited presenters on the major topics. Each talk will consist of 20 minutes except for invited talks where 30 minutes will be allocated. The poster session will compliment the oral program and offers a venue for innovative research on this subject to be presented and discussed.

continued on page 15

24th International Applied Geochemistry Symposium... continued from page 14

SPECIAL SESSIONS

New Frontiers for Exploration in Glaciated Terrain

CoChairs: Beth McClenaghan (Geological Survey of Canada), Roger Paulen (Alberta Geological Survey), Bill Coker and Chris Benn (BHP Billiton)

Since the discovery of economic diamond deposits in northern Canada in the 1990s, the application of indicator mineral methods has been expanding to include a broader spectrum of commodities including base metals and uranium. The application of till geochemistry continues to be an important part of the drift exploration methodology. The session will highlight recent developments in indicator mineral and till geochemistry methods as well as recent case histories for diamonds, base metals, gold, PGE and uranium.

Applied Geochemistry of Geological Storage of CO2

Cochairs: William D. Gunter & Ernie H. Perkins (Honorary Cochair: Brian Hitchon)

The United States, the European Union, Australia and Canada are all pursuing Carbon Capture and Storage (CCS) demonstration projects as a option for reducing GHG emissions from usage of fossil fuel at large Final Emitters (LFEs). Currently in Canada, CCS is being considered by Alberta, Saskatchewan and Nova Scotia. A critical part of CCS is the geological storage of the CO₂. This involves finding a secure geological site in depleted oil and gas reservoirs or deep saline aquifers. An unusual facet of geological storage is that the containment of the CO₂ in

the storage reservoir becomes more secure with time due to geochemical reactions where the CO₂ may potentially dissolve in the formation water and react with the formation minerals neutralizing the carbonic acid and precipitating carbonate minerals. There are two time scales involved, (i) the short operational time scale (10s of years) when CO₂ is being actively injected into the reservoir and (ii) the long geochemical time scale, post closure after injection has ceased, when water – rock reactions are most active (100s to 1000s of years). Verification of geochemical trapping of CO₂ is difficult given the long time scales involved. This session will be devoted to addressing this issue through experimental, natural analogues, industrial analogues and numerical simulation.

Applied Geomicrobiology: New Frontiers in Exploration and **Environmental Geomicrobiology**

Cochairs: Christopher Weisener (U. Windsor), David A. Fowle (U. Kansas)

This forum will address aspects of applied geomicrobiology, specifically the role of microbes on elemental cycling in near-surface environments. The focus of the session will include topics related to: microbial transformations in mine settings, metal sequestration and mobilization associated with ore deposits, bioremediation and augmentation, isotopic, molecular and new methods in applied geomicrobiology.

continued on page 16

Exploration & Mining Geochemistry

New International Workshop for 2008

Half Day Modules Full Immersion

Presenters include: Prof. Kurt Kyser **Colin Dunn Dave Lawie Gwendy Hall** Mike Whitbread **Steve Sugden Dennis Arne**

...more to be announced

Geochemistry Essentials

Surface Sampling Strategies for Exploration

Geoanalysis, including Partial Extractions & Instrumentation

Introduction to Exploratory Data Analysis

QC From Exploration to Production

Lithogeochemistry for Mapping & Quantification of Alteration

Phase-specific & Real-time Assaying Methods

Advanced Exploratory Data Analysis

Adding Value to Exploration with Isotopes & New Techniques **Biogeochemistry**

Commodity Sessions – Au,Cu,Mo; Ni,PGM; U; Base metals (Cu,Pb,Zn)

For more information visit www.ioglobal.net or email workshop@ioglobal.net

Places strictly limited and offered on a first come - first served basis; register early to avoid disappointment

Perth • Melbourne • Brisbane

PAGE 16 NUMBER 140 EXPLORE

24th International Applied Geochemistry Symposium... continued from page 14

Mine Waste Characterization: State of the Practice, Problems, and Future Directions

Cochairs: David Bird (Colorado Division of Reclamation, Mining & Safety, DNR), James Ranville, (Colorado School of Mines), Ronald Schmiermund (Economic & Environmental Geochemistry Inc.)

Predicting long-term environmental impacts from mining wastes at hard rock metal mining sites is an increasingly necessary component of the permitting and financing processes. In addition, there are escalating demands for minimal long-term water treatment, environmentally-acceptable pit lakes and chemically stable waste dumps and tailings storage facilities. The current boom is putting intense pressure on the industry to meet these challenges and the consequences of failure are becoming increasingly high-profile with attendant global corporate liabilities. Unfortunately, accurate predictions of waste behavior have generally been difficult to obtain, and ideal "walk-away" closures in most cases remain elusive.

The understanding of acid rock drainage and associated processes must be considered relatively sophisticated, albeit incomplete. However, understanding appears to be disproportionately great relative to success in prediction and prevention. In part, this under-achievement stems from a failure to thoroughly integrate, in a timely fashion, waste characterization programs with economic geology, environmental baselines, mine planning and development in addition to the prevailing regulatory requirements. In part, the shortcomings of the available predictive tools and techniques are contributors to the problem.

This session will explore the state of the practice of mine waste characterization and its place in the overall creation of a modern mine. Lessons learned from the past will be considered, and how those lessons can be applied to improving methodologies, and the interpretation of the data derived from the existing methodologies. Papers will be sought in the following areas: Evaluations of overall success/failure of previous efforts, Corporate programs and perspectives for integrating waste characterization, Multinational case studies of projects with lessons learned, State-of-the art analytical, simulation and modeling tools.

NORTH ATLANTIC MINERALS SYMPOSIUM (6th)

Cochairs: Andrew Kerr (Geological Survey of Newfoundland and Labrador), Gerry Stanley (Geological Survey of Ireland), Lawrence Winter (Altius Minerals Corporation)

The NAMS concept is intended to bring together representatives of industry, government geoscience, and university research who share a common interest in the geology and mineral potential of the North Atlantic region. This focus is defined loosely to include eastern North America and western Europe, adjacent northern lands, and Atlantic islands. NAMS seeks reviews and ideas connected to the regional and district metallogeny of these regions, exploration and research case studies of all types, and also wider conceptual contributions related to metallogeny that are applicable in the region. In the wider context of this conference, contributions related to the use of applied

geochemical techniques in the North Atlantic exploration environment are also welcome. The North Atlantic Minerals Symposium (NAMS) is a joint initiative of the Geological Survey of Ireland and the Geological Survey of Newfoundland and Labrador (Canada).

GENERAL SESSIONS

- Classic mining districts: exploration and environmental geochemistry in the shadow of headframes
- Mapping Geochemical Data
- Geochemistry in the industry sector: exploration and environmental applications.
- Recent & Classic Exploration Case Studies
- Ore Deposit-Forming Systems: A Geochemical Perspective
- Biogeochemistry
- Data management, Statistical Analysis, and Interpretation
- Lithogeochemical Applications
- Analytical Geochemistry: rocks, regolith, soils, organics, water, and air
- Geochemistry in the Government Sector: exploration and environmental applications
- Geochemistry and Health
- Geochemical Aspects of Mine Waste
- Applied Mineral-Chemical Interactions
- New technologies in Geochemical Exploration to Waste Management
- Remote Sensing and Petrophysics

POSTERS

The Local Organizing Committee will supply a poster boards (dimensions to follow shortly). Velcro will be provided to attach posters. The authors should also prepare some 11" by 17" or 8.5" by 14" copies of their poster to hand out to interested delegates. On Tuesday afternoon, authors will have the opportunity to present (4 slide ppt) for 5 minutes on their topic.

For the convenience of the delegates, posters may be printed by the Organization at a cost of \$125 Cdn. Those delegates interested should send to the Congress secretariat before May 1st, 2009 in a pdf (containing a page with the final dimensions, within the established limits) and it will be printed and placed on the Congress boards.

SUBMISSION OF EXTENDED ABSTRACTS AND PAPERS

Extended abstracts should be sent to dlentz@unb.ca as a single e-mail attachment, saved as a Microsoft Word document, using the extended abstract template, with figures and tables embedded in the document. Format guidelines for the abstracts are available at *Geochemistry: Exploration, Environment, Analysis*" (GEEA) "Instructions to Authors", at the website listed below.

http://www.geolsoc.org.uk/template.

cfm?name=journals_geea_home_page

Papers from the conference proceedings will be published in "Geochemistry: Exploration, Environment, Analysis" (GEEA), produced jointly by the Association of Applied

24th International Applied Geochemistry Symposium... continued from page 16

Geochemists and the Geological Society of London. Manuscripts, indicating '24th IAGS', should be submitted by September 30, 2009 to: Marcia Scrimgeour, Editorial Office Manager, *Geochemistry: Exploration, Environment, Analysis*, P.O. Box 26099, 72 Robertson Road, Nepean, Ontario, K2H 9R0, CANADA. Email: geea@sympatico.ca

STUDENT BURSARIES FOR TRAVEL

A limited number of bursaries will be available to students to assist with travel to the conference. Information and application forms will be posted on the symposium website shortly.

IMPORTANT DATES

- January 15, 2009: Start of submission of Extended Abstracts
- February 15, 2009: Deadline for submission of Extended Abstracts
- March 25, 2009: Deadline for student bursary applications
- April 10, 2009: Deadline to inform the students if their applications to the Student Travel Bursary have been successful
- April 2nd, 2009: Deadline to inform authors of their acceptance (oral or poster)
- April 30, 2009: Deadline for Early-bird registration, a surcharge of 50 will be levied thereafter, and field trips will be decided. All authors of extended abstracts must register by this date, or they will be removed from the program.
- September 30, 2009: Deadline for submission of papers to Geochemistry: Exploration, Environment, Analysis for the Proceedings Special Issue

CONTACT US

For further Information: Department of Geology 2 Bailey Drive, Box 4400 University of New Brunswick Fredericton, NB E3B 5A3 CANADA Tel: 1-506-453-4804; FAX: 1-506-453-5055

email: geology@unb.ca

The Department of Geology at the University of New Brunswick is active in both teaching and research in New Brunswick and around the world and has a long and proud history with strong undergraduate and graduate programs. Website: http://www.unb.ca/fredericton/science/geology/

LOCAL ORGANIZING COMMITTEE

Chairman, Prof. David Lentz, University of New Brunswick (UNB), Fredericton, NB., Canada

email: dlentz@unb.ca

Honorary Chairman: Prof. Gerry Govett

Treasurer: Dr. Steven McCutcheon (NBDNR-Minerals,

UNB Adjunct Prof.)

Technical Program Co-chairs: Prof. David Keighley (UNB) & Dr. Kay Thorne (NBDNR-Minerals)

Field Trip Chair: Dr. James Walker (NBDNR-Minerals) **Workshops:** Dr. Don Fox (NB DOE) & Terry Goodwin (NS DNR)

Exhibits: Dr. Nick Susak (UNB), Dr. Doug Hall (UNB) **Sponsorships:** Ross Gilders (NB Research & Productivity Council)

UNB Geology Administrator: Tammy O'Donnell Guest Program: Christine Lodge (UNB Geology)
Social Program: Director of UNB Conference Services & Destination Sales Coordinator, Rendez-Vous Fredericton Publicity: Dr. David Lentz

The Association of Applied Geochemists

announces the



2008 AAG
Student
Paper
Competition

The AAG is calling for nominations for the 17th biennial Student Paper Competition. The paper must address an aspect of exploration geochemistry or environmental geochemistry related to mineral exploration and represent research performed as a student. The student must be the principal author and the paper must have been published in *Geochemistry: Exploration, Environment, Analysis* no more than three years after completion of the degree. A nomination may be made by anyone familiar with the work of the student.

Deadline for receipt of nominations is December 31, 2008.

The winner will receive:

A cash prize of \$1000CAD generously donated by SGS Minerals Services.

A 2-year membership of AAG, including the society's journal (GEEA), EXPLORE newsletter, publication of an abstract and CV of the winner, a certificate of recognition and \$500US towards expenses to attend an AAG-sponsored meeting, courtesy of AAG.

Nominations and a digital copy of the paper should be sent to:

Dr David Cohen
Chair, Student Paper Competition
School of BEES
The University of New South Wales
UNSW NSW 2052
Australia
Email: d.cohen@unsw.edu.au

The results of the 2008 competition will be announced at the 24th IAGS in mid 2009.

Further details are available from the chair of the committee or the AAG Students' page at http://www.appliedgeochemists.org/

PAGE 18 NUMBER 140 EXPLORE



CALENDAR OF EVENTS

International, national, and regional meetings of interest to colleagues working in exploration, environmental and other areas of applied geochemistry. These events also appear on the AAG web page at: www.appliedgeochemists.org

2008

- September 8-10, 2008, 9th International Congress for Applied Mineralogy, Brisbane, Australia. Website: http://www/icam2008.com
- September 14–18, 2008, 5th International Conference on Uranium Mining and Hydrogeology. Freiberg, Germany. Website: URL: http://www.geo.tu-freiberg.de/umh
- September 15-20, 2008. Association of Environmental and Engineering Geologists 51st Annual Meeting, New Orleans, Louisiana USA. Website: http://www.aegweb.org/i4a/pages/index.cfm?pageid=3836
- October 2-3, 2008. 2008 NGWA/U.S. EPA Remediation of Abandoned Mine Lands Conference. Denver, USA.
 Website: www.ngwa.org/DEVELOPMENT/conferences/ details/08100025109.aspx
- October 5-8, 2008. Geological Society of America Annual Meeting, Houston, Texas, USA. Website: www. geosociety.org/meetings/index.htm
- October 27-29. 2008. Exploration on the Edge. ARC Centre of Excellence in Ore Deposits (CODES), Hobart, Tasmania.

Website: http://fcms.its.utas.edu.au/scieng/codes/index.asp

- October 27-November 7, 2008. Ore Deposit Models and Exploration Strategies. ARC Centre of Excellence in Ore Deposits (CODES), Hobart, Tasmania.
 Website: http://fcms.its.utas.edu.au/scieng/codes/cpage. asp?lCpageID=55
- November 4–6 2008. 22nd Colloquium of African Geology and 13th Conference of the Geological Society of Africa. Hammamet, Tunisia. E-mail: afric2008@gmail. com
- November 14-20, 2008, Exploration & Mining Geochemistry International Workshop, Perth, Western Australia. Website www.ioglobal.net.
- December 3-12, 2008. Modular Course in Exploration Geochemistry. Sudbury, Ontario, Canada. Email: spiercey@laurentian.ca. Website: http://earthsciences.laurentian.ca.
- December 15-19, 2008. American Geophysical Union Fall Meeting. San Francisco, USA. Website: www.agu.org/ meetings/fm08/

2009

- January 26-29, 2009 Mineral Exploration Roundup 2009.
 Vancouver, B.C. Canada. Website: http://www.amebc.ca/roundupoverview.htm
- March, 2009. Ore Deposits of South America. ARC Centre of Excellence in Ore Deposits (CODES), Hobart, Tasmania.

Website: http://www.fcms.its.utas.edu.au/scieng/codes/cpage.asp?lCpageID=55

continued on page 19



Association of Applied Geochemists web site:

appliedgeochemists.org



CALENDAR OF EVENTS

continued from page 18

- March 1-4, 2009. Prospectors and Developers Association of Canada Annual Convention, Toronto, Canada. Website: http://http://www.pdac.ca
- May 24 27, 2009. Geological Association of Canada/ Mineralogical Association of Canada Annual Meeting Toronto, Canada.

Website: http://www.gac.ca/activities/index.php

- June, 2009. Ore Deposit Geochemistry, Hydrology and Geochronology. ARC Centre of Excellence in Ore Deposits (CODES), Hobart, Tasmania.
 Website: http://www.fcms.its.utas.edu.au/scieng/codes/ cpage.asp?lCpageID=55
- June 1 4, 2009. 24th International Applied Geochemistry Symposium, Fredericton, New Brunswick, Canada Website:http://www.unb.ca/conferences/IAGS2009
- June 22 26, 2009. Goldschmidt 2009. Davos, Switzerland. Website: http://www.goldschmidt2009.org
- August 17-20, 2009. Society for Geology Applied to Mineral Deposits 10th Biennial Meeting, Townsville, Australia. Email: SGA2009@jcu.edu.au
- September 7-11, 2009. Geoanalysis 2009. Drakensberg Region, South Africa.
 Website: http://geoanalysis2009.org.za
- September 21-26, 2009. Association of Environmental and Engineering Geologists 52nd ANNUAL MEETING, Lake Tahoe, USA. Website: http://www.aegweb.org/i4a/pages/ Index.cfm?pageID=3696
- October 18-21, 2009. Geological Society of America Annual Meeting. Portland, Oregon, USA. Website: www. geosociety.org/meetings/index.htm

2010

- April, 2010. 27th Society for Environmental Geochemistry and Health, European Conference, Galway, Ireland. Website: http://www.nuigalway.ie/segh2010/
- May, 2010. Geological Association of Canada/ Mineralogical Association of Canada Annual Meeting Calgary, Canada. Website: http://www.gac.ca/activities/ index.php
- July 14-18, 2010. Goldschmidt 2010. Knoxville, USA. Website: www.geochemsoc.org/news/conferencelinks/

 October 31-November 3, 2010. Geological Society of America Annual Meeting. Denver, Colorado, USA. Website: www.geosociety.org/meetings/index.htm

2012

- August 5–15, 2012. 34th International Geological Congress, Brisbane, Australia. Website: http://www.ga.gov.au/igc2012
- 2012. Geoanalysis 2012. Brazil

Please let this column know of your events by sending details to:

Beth McClenaghan

Geological Survey of Canada 601 Booth Street Ottawa, Ontario, CANADA K1A 0E8 Email: bmccmwba@nrcan.gc.ca 613-992-7805



Mineral Exploration Roundup 2009 Short Course

January 26 to January 29, 2009 Vancouver, British Columbia, CANADA Website: http://www.amebc.ca/default.htm

Sources and Sinks in Hydrothermal Systems

Presented by AME BC, Dick Tosdal, Stephen Cox, Mike Lesher, Kurt Kyser, Peter Hollings, Wayne Goodfellow Date: January 24-25, 2009

Overview:

Ore deposits require a source of fluids, ligands, and metals as well as a transport system and sink for precipitation in economic concentration. These involve the interplay between magmatic activity, basin development, and tectonics. Understanding the scale of those systems and recognizing where within a paleohydrothermal system on might be can provide important vectors toward undiscovered resources.

This 2-day course will examine the sources and sinks of hydrothermal deposits. The course is aimed at mineral exploration, government, academic and student geologists, and provides an opportunity to meet and exchange data and views with leading researchers in the field.

PAGE 20 NUMBER 140 EXPLORE



RECENT PAPERS

This list comprises titles that have appeared in major publications since the compilation in **EXPLORE** Number 139. Journals routinely covered and abbreviations used are as follows: Economic Geology (EG); Geochimica et Cosmochimica Acta (GCA); the USGS Circular (USGS Cir); and Open File Report (USGS OFR); Geological Survey of Canada papers (GSC paper) and Open File Report (GSC OFR); Bulletin of the Canadian Institute of Mining and Metallurgy (CIM Bull.): Transactions of Institute of Mining and Metallurgy, Section B: Applied Earth Sciences (Trans. IMM). Publications less frequently cited are identified in full. Compiled by L. Graham Closs, Department of Geology and Geological Engineering, Colorado School of Mines, Golden, CO 80401-1887, Chairman AEG Bibliography Committee. Please send new references to Dr. Closs, not to EXPLORE.

Albanese, S., 2008. Evaluation of the bioavailability of potentially harmful elements in urban soils through ammonium acetate-EDTA extraction: a case study in southern Italy. <u>in</u> Geochemistry: Exploration, Environment, Analysis: <u>8</u>(1): 49-57.

Arcos, D., et al., 2008. Geochemical modeling of the weathering zone of the "Mina Fe" U deposit Spain: A natural analogue for unclear spent fuel alteration and stability processes in radwaste disposal. Applied Geochem. 23(4): 807-821.

Armienta, M.A., et al., 2008. Arsenic distribution in mesquite (Prosopis laevigata) and huizache (Acarcia farnesiana) in the Zimapan mining area, Mexico. <u>in</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(2): 191-198.

Modular Course in Exploration Geochemistry

Date: December 3-12, 2008.

Location:

Laurentian University, Sudbury, Ontario, Canada.
Information: Dr. Steve Piercey,
Mineral Exploration Research Centre,
Department of Earth Sciences,
Laurentian University,
Willet Green Miller Centre,
933 Ramsey Lake Road,
Sudbury, ON, Canada, P3E 2C6;
tel. +1.705.675.1151 x4595; fax +1.705.675.4898;

e-mail: spiercey@laurentian.ca; website: http://earthsciences.laurentian.ca

Astrom, M.E., et al., 2008. Niobium in boreal stream waters and brackish-water sediments. <u>in</u> (2): 139-148.

Ayuso, R.A. and Foley, N.K., 2008. Anthropogenic and natural lead isotopes in Fe-hydroxides and Fe-sulphates in a watershed associated with arsenic-enriched groundwater, Maine, USA. <u>In</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(1): 77-89.

Barrett, T.J., Dawson, G.L., and MacLean, W.H., 2008. Volcanic Stratigraphy, Alteration, and Sea-Floor Setting of the Paleozoic Feitas Massive Sulfide Deposit, Aljustrel, Portugal. EG <u>103</u>(1): 215-239.

Bettiol, C., et al., 2008. Evaluation of microwave-assisted acid extraction procedures for the determination of metal content and potential bioavailability in sediments. Applied Geochem. 23(5): 1140-1151.

Bissig, T., et al., 2007. Vein carbonates in the low sufidation epithermal Au-AG district of El Penon, II Region, Chile. Revista Geol de Chile 34(2): 291-

Bradshaw, G.D., et al., 2008. Genesis of the Wolverine Volcanic Sediment-Hosted Massive Sulfide Deposit, Finlayson Lake District, Yukon, Canada: Mineralogical, Mineral Chemical, Fluid Inclusion, and Sulfur Isotope Evidence. EG 103(1): 35-60.

Budd, A.R. and Skirrow, R.G., 2007. The Nature and Origin of Gold Deposits of the Tarcoola Goldfield and Implications for the Central Gawler Gold Province, Australia. EG <u>102</u>(8): 1541-1563.

Bulter, B.A., Ranville, J.F., and Ross, P.E., 2008. Direct versus indirect determination of suspended sediment associated metals in a mining-influenced watershed. Applied Geochem. <u>23</u>(5): 1218-1231.

Cabral, A.R., et al., 2007. Supergene leaching and formation of platinum in alluvium: Evidence from Serro, Minas Gerais, Brazil. Min. Petrol. 90(1/2): 141-

Canovas, C.R., et al., 2007. Hydrogeochemical characteristics of the Tinto and Odial Rivers (SW Spain). Factors controlling metal contents. Sci. Total Environ. 373(1): 363-

Chi, G. et al., 2008. Geologic, Geochemical, and Geochronological Constraints on the Genesis of Gold Mineralization at Poplar Mountain, Western New Brunswick, Canada. Explor. Min. Geol. <u>17</u>(1/2): 101-130.

Chiaradia, M., et al., 2008. Geologic Setting, Mineralogy, and Geochemistry of the Early Tertiary Au-Rich Volcanic-Hosted Massive Sulfide Deposit of La Plata, Western Cordillera, Ecuador. EG 103(1): 161-183.



RECENT PAPERS

continued from Page 20

Chopin, E.I.B. and Alloway, B.J., 2007. Trace element partitioning and soil particle characterization around mining and smelting areas at Tharsis, Riotinto, and Huelva, SW Spain. Sci. Total Environ. 373(2/3): 488.

Cicchella, D., et al., 2008. Urban geochemical mapping in Campania region (Italy). <u>in</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(1): 19-29.

Cicchella, D., et al., 2008. Platinum group element distribution in the soils from urban areas in the Campania Region (Italy). -n Geochemistry: Exploration, Environment, Analysis 8(1): 31-40.

Cicchella, D., et al., 2008. Heavy metal pollution and Pb isotopes in urban soils in Napoli, Italy. <u>in</u> Geochemistry: Exploration, Environment, Analysis 8(1): 103-112.

Colombo, C., et al., 2008. Complexation of platinum, palladium, and rhodium with organic ligands in the environment. <u>In</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(1): 91-101.

Costagliola, P., et al., 2008. Impact of ancient metal smelting on arsenic pollution in the Pecora River Valley, Southern Tuscany, Italy. Applied Geochem. <u>23</u>(5): 1241-1259.

Creighton, S., et al., 2008. Diamondiferous peridotic microxenoliths from the Diavik Diamond Mine. NT Contrib. Min. Petrol. <u>155(5)</u>: 541-554.

Daliran, F., 2008. The carbonate rock-hosted epithermal gold deposit of Agdarreh, Takab geothermal field, NW Iran – hydrothermal alteration and mineralization. Min. Deposita 43(4): 383-404.

Das, A.K., et al., 2007. A review on molybdenum determination in solid geological samples. Talanta <u>71(3)</u>: 987.

Del Rio Salas, R., et al., 2008. Geology, Geochemistry, and Re-Os systematics of manganese deposits from the Santa Rosalia Basin and adjacent areas of Baja California Sur, Mexico. Min. Deposits 43(4): 467-482.

Dewing, K., 2007. Are Your Data Good Enough: A Checklist for Mining Prospects. Geoscience Canada <u>34(2)</u>: 65-69.

Dickson, B.L. and Giblin, A.M., 2007. Effective exploration for uranium in South Australia paleochannels. Trans. IMM <u>116(2)</u>: B50-

Eapala, M.P., Parry, D., and Noller, B., 2007. Dynamics of arsenic in the mining sites of Pine Creek Geosyncline, Northern Australia.

Faure, G. and Mensing, T.M., 2007. Introduction to Planetary Science: The Geological Perspective. Springer. 526 p.

Fedele, L., et al., 2008. The rare earth element distribution over Europe: geogenic and anthropogenic sources. <u>in</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(1): 3-18.

Fernandez-Caliani, J.C., Borga-Brioso, and Perez-Lopez, R., 2008. Long-term interaction of wollastinite with acid mine water and effects on arsenic and metal content. Applied Geochem. 23(5): 1288-1289.

Foley, N.K. and Ayuso, R.A., 2008. Mineral sources and transport pathways for arsenic release in a coastal watershed, USA. <u>in</u> Geochemistry: Exploration, Environment, Analysis 8(1): 59-75.

Fonseca, R.O.C., et al., 2007. How chalcophile is rhenium? An experimental study of the solubility of Re in sulphide mattes. Earth Planet. Sci. Letters <u>260</u>(3/4): 537-

Frau, F., Biddau, R., and Fanfani, L., 2008. Effect of major anions on arsenate desorption from ferrihydrite-bearing natural samples. Applied Geochem. 23(6): 1451-1466.

Gandy, C.J., Smith, J.W.N., and Jarris, A.P., 2007. Alternation of mining-derived pollutants in the hyporheic zone: A review. Sci. Total Environ. 373(2/3): 435-

Garrett, R.G., et al., 2008. Variations in geochemical background levels for Jamaican soils. <u>in</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(2): 149-156.

Garrett, R.G., et al., 2008. The presence of anomalous trace element levels in present day Jamaican soils and the geochemistry of Late-Miocene or Pliocene phosphorites. Applied Geochem. 23(4): 822-834.

Gemmell, J.B., 2007. Hydrothermal Alteration Associated with the Gosowong Epithermal Au-Ag Deposit, Halmahera, Indonesia: Mineralogy, Geochemistry, and Exploration Implications. EG 102(5): 893-922.

Groat, L.A. (ed.), 2007. Geology of Gem Deposits. Min. Assoc. Canada Short Course V. 37. 288 p.

Gurvich, E.G., 2006. Review of Metalliferous Sediments of the World – Fundamental Theory of Deep-Sea Hydrothermal Sedimentation. Springer. 416 p.

Haffert, L. and Craw, D., 2008. Mineralogical controls on environmental mobility of arsenic from historic mine processing residues, New Zealand. Applied Geochem. <u>23</u>(6): 1467-1483.

PAGE 22 NUMBER 140 EXPLORE



RECENT PAPERS

continued from Page 21

Haldar, S.K., 2007. Exploration Modeling of Base Metal Deposits. Elsevier. 227 p.

Han, Y., et al., 2008. Particulate-associated potentially harmful elements in urban road dusts in Xi'an, China. Applied Geochem. <u>23</u>(4): 835-845.

Haus, K.L., et al., 2008. Analysis of arsenic speciation in mine contaminated lacustrine sediment using selective sequential extraction, HR-ICPMS and TEM. Applied Geochem. 23(4): 692-704.

Hoel, K.O., 2008. Getting the Geo into Geomet. SEG Newsletter 73(1): 11-15.

Holl, R., Kling, M., and Schroll, E., 2007. Metallogenesis of germanium – A Review. Ore Geol. Rev. 30(3/4): 145-

Hough, R.M., et al., 2007. Gold nuggets: Supergene or hypergene? Aust. J. Earth Sci. <u>54</u>(7): 959.

Hough, R.M., et al., 2008. Naturally occurring gold nanoparticles and nanoplates. Geology <u>36</u>(7): 571-574.

Izbilki, J.A., et al., 2008. Chromium, chromium isotopes, and selected elements, western Mojave Desert, USA. Applied Geochem. 23(5): 1325-2352.

Jefferson, C.W. and Delaney G. (Eds), 2007. EXTECH IV: Geology and Uranium Exploration. Technology of the Proterozoic Athabaska Basin, Saskatchewan and Alberta. Geol. Surv. Can. Bull 588. 645 p.

Juhas, A. and Rice, L.R., 2008. The Exploration Mine Development Process: A Modern Application of the Scientific Method. The Professional Geologist <u>45</u>(3): 60-61.

Kahmann, J.A., Seaman, J., III, and Zhou, M.F., 2008. Evaluating Trace Elements as Paleoclimate Indicators: Multivariate Analysis of Late Mississippian Pennington Formation Paleosols, Kentucky, USA. J. Geol. <u>116</u>(3): 254-268.

Keegan, E., et al., 2008. The provenance of Australian uranium ore concentrates by elemental and isotopic analysis. Applied Geochem. <u>23</u>(4): 765-777.

Kohfahl, C., et al., 2008. Recharge sources and hydrogeochemical evolution of groundwater in semi-arid and karstic environments: A field study in the Granada Basin (Southern Spain). Applied Geochem. <u>23</u>(4): 846-862.

Konhauser, K., 2007. Introduction to Geomicrobiology. Blackwell Science. 425 p.

Krysiak, A. and Karczweka, A., 2007. Acid extractability in soils of former arsenic mining and smelting, SW Poland. Sci. Total Environ. 379(2/3): 100-

Layton-Matthews, D., et al., 2008. Distribution, Mineralogy, and Geochemistry of Selenium in Felsic Volcanic-Hosted Massive Sulfide Deposits of the Finlayson Lake District, Yukon Territory, Canada. EG 103(1): 61-88.

Li, J., et al., 2007. A simplified method for estimation of jarosite and acid-forming sulfates in acid mine wastes. Sci. Total Environ. 373(1): 391.

Liebscher, A. and Heinrich, C.A. (Eds), 2007. Fluid-Fluid Interactions. Min. Soc. Am. Rev. Min. Geochem. V. 65. 432 p.

Le Gleuher, M., et al., 2008. Mineral hosts for gold and trace metals in regolith at Boddington gold deposit and Scuddler massive copper-zinc sulphide deposit, Western Australia: An LA-ICP-MS study. in Geochemistry: Exploration, Environment, Analysis 8(2): 157-172.

Lima, A., et al., 2008. Interpolation methods from geochemical maps: a comparative study using arsenic data from European stream waters. <u>in</u> Geochemistry: Exploration, Environment, Analysis 8(1): 41-48.

Lottermoser, B.G., Ashley, P.M., and Munksgaard, N.C., 2008. Biogeochemistry of Pb-Zn gossans, northwest Queensland, Australia: Implications for mineral exploration and mine site rehabilitation. Applied Geochem. <u>23</u>(4): 723-742.

Makkonen, H.V., Makinen, J., and Kontoniemi, O., 2008. Geochemical discrimination between barren and mineralized intrusions in the Svecofennian (1.9 Ga) Kotalahti Nickel Belt, Finland. Ore Geol. Rev. 33(1): 101.

Malmstrom, M.E., Berglund, S., and Jarsjo, J., 2008. Combined effects of spatially variable flow and mineralogy on the attenuation of acid mine drainage in groundwaters. Applied Geochem. <u>23(6)</u>: 1419-1436.

Mamaeva, E.I., 2006. Tract elements as indicators of conditions of the ore-forming mineralization in rocks of Gulinsky massif. Proc. Russian Min. Soc. 135(6): 54.-

Martin, F., et al., 2008. Soil alteration by continued oxidation of pyrite tailings. Applied Geochem. <u>23(5)</u>: 1152-1165.

Martinez, J., et al., 2007. Determination of the geochemical background in a metal mining site: Example of the mining district of Linares (south Spain). J. Geochem. Explor. 94(1-3): 19-



RECENT PAPERS

continued from Page 22

Mauk, J.L. and Simpson, M.P., 2007. Geochemistry and Stable Isotope Composition of Altered Rock at the Golden Cross Epithermal Au-Ag Deposit, New Zealand. EG <u>102(5)</u>: 841-871.

Maxwell, P. and Guj, P. (Eds), 2006. Australian Mineral Economics. Aus. IMM Mono 24. 240 p.

McInnis, M., et al., 2008. Trace elements in native gold by solution ICP-MS and their use in mineral exploration: A British Columbia examples. Applied Geochem. <u>23(5)</u>: 1076-

McLeod, M.J., Hoy, D., and Thorne, K.G., 2008. History, Tectonic Setting, and Models for Intrusion-Related Gold Deposits in Southwestern New Brunswick, Canada: Examples from the Clarence Stream Area. Explor. Min. Geol. <u>17</u>(1/2): 1-12.

Moldovan, B.J., Hendry, M.J., and Harrington, G.A., 2008. The arsenic source term for an in-pit uranium mine tailings facility and its long-term impact on the regional groundwater. Applied Geochem. 23(6): 1437-1450.

Mukherjee, A. and Fryer, A.E., 2008. Deeper groundwater chemistry and geochemical modeling of the arsenic affected western Bengal basin, West Bengal, India. Applied Geochem. 23(4): 863-894.

Mungall, J.E., 2007. Crystallization of magmatic sulfides: An empirical model and application to Sudbury ores. GCA <u>71</u>(11): 2809.

Naftz, D., et al., 2008. Anthropogenic influences on the input and biogeochemical cycling of nutrients and mercury in Great Salt Lake, Utah, USA. Applied Geochem. <u>23(6)</u>: 1731-1744.

Navarro, A., et al., 2008. Metal mobilization from basemetal smelting slag dumps in Sierra Almagrera (Almeria Spain). Applied Geochem. <u>23</u>(4): 895-913.

Neaman, A., et al., 2008. Trace element associations with Fe- and Mn-oxides in soil nodules: Comparison of selective dissolution with electron probe microanalysis. Applied Geochem. 23(4): 778-782.

Nutt, C.J. and Hofstra, A.H., 2007. Bald Mountain Gold Mining District, Nevada: A Jurassic Reduced Intrusion-Related Gold System. EG <u>102(6)</u>: 1129-1155.

Oyarzun, R., et al., 2007. Mineral deposits and Cu-Zn-As dispersion-contamination in stream sediments from the semiarid Coquimbo Region, Chile. Environ. Geol. <u>53(2)</u>: 283.

Parello, A., et al., 2008. Geochemical characterization of surface waters and groundwater resources in the Managua area (Nicaragua, Central America). Applied Geochem. 23(4): 914-931.

Peng, B., et al., 2007. Mineralogical and geochemical constraints on environmental impacts from waste rock at Taujiang Mu-ore deposit, central Hunan, China. Environ. Geol. <u>52</u>(7): 1127.

Piercey, S.J., et al., 2008. Petrology and U-Pb Geochronology of Footwall Porphyritic Rhyolites from the Wolverine Volcanogenic Massive Sulfide Deposit, Yukon, Canada: Implications for the Genesis of Massive Sulfide Deposits in Continental Margin Environments. EG <u>103</u>(1): 5-33.

Pina, R., et al., 2008. Mineralogy and geochemistry of platinum-group elements in the Aquablanca Ni-Cu deposit (SW Spain). Min. Petrol. <u>92</u>(1): 259-282.

Ravenelle, J.-F., Lutes, G.G., and Hynes, A.J., 2008. Architecture of Gold Mineralization at Anomaly A of the Clarence Stream Deposits, Southern New Brunswick. Explor. Min. Geol. 17(1/2): 85-100.

Reith, F. and McPhail, D.C., 2007. Mobility and microbially mediated mobilization of gold and arsenic in soils from two gold mines in semi-arid and tropical Australia. GCA <u>71(5)</u>: 1183.

Resing, J.A., et al., 2007. Venting of Arid-Sulfate Fluids in a High-Sulfidation Setting at NW Rota-1 Submarine Volcano on the Mariana Arc. EG <u>102</u>(6): 1047-1061.

Rusk, B.G., Lowers, H.A. and Reed, M.H., 2008. Trace elements in hydrothermal quartz: relationships to cathodoluminescent textures and insights into vein formation. Geology 36(7): 547-550.

Salvarredy-Aranguren, M.M., et al., 2008. Contamination of surface waters by mining waste in the Milluni Valley (Cordillera Real, Bolivia): Mineralogical and hydrological influences. Applied Geochem. <u>23</u>(5): 1299-1324.

Samal, A.R., 2008. Digital Mineral Resource Modeling: An Overview. The Professional Geologist 45(3): 62-63.

Sanchez-Espano, J., et al., 2008. The acid mine pit lakes of the Iberian Pyrite Belt: An approach to their physical liminology and hydrogeochemistry. Applied Geochem. 23(5): 1260-1287.

Sasaki, K., et al., 2008. Immobilization of Se (VI) in mine drainage by permeable reactive barriers: column performance. Applied Geochem. 23(5): 1012-1022.

PAGE 24 NUMBER 140 EXPLORE



RECENT PAPERS

continued from Page 23

Sarala, R.H., 2007. Distinguishing glaciogenic deposits in southern Finnish Lapland: Implications for exploration. Trans. IMM <u>116(1)</u>: B22-

Sillitoe, R.H., 2007. Hypogene Reinterpretation of Supergene Silver Enrichment at Chanarcillo, Northern Chile. EG <u>102</u>(5): 777-781.

Silver, D.B., 2008. Super Cycle: past, present, and future (part 1). Min. Eng. <u>60</u>(5): 72-77.

Somarin, A.K. and Lentz, D.R., 2008. Mineralogy, geochemistry and fluid evolution of a fossil hydrothermal system in the Paleogene Mendejin volcanic sequence, East Azarbaijan, Iran. Min. Petrol. Springer.

Staff, USGS Mineral Information Team, 2008. Mining Review. Min. Eng. <u>60</u>(5): 31-43.

Stanley, C.R., 2008. Missed hits or near misses: determining how many samples are necessary to confidently detect nugget-borne mineralization. <u>in</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(2): 129-138.

Stanley, C.R. and Lawie, D., 2008. Thompson-Howarth error analysis: unbiased alternatives to the large sample method for assessing non-normally distributed measurement errors in geochemical samples. <u>in</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(2): 173-182.

Stanley, C.R. and Noble, R.R.P., 2008. Quantitative assessment of the success of geochemical exploration techniques using minimum probability methods. <u>in</u> Geochemistry: Exploration, Environment, Analysis <u>8</u>(2): 115-127.



Association of Applied Geochemists Student Membership \$10 US

Encourage a student to join!

Taylor, C.D., et al., 2008. The Metallogeny of Late Triassic Rifting of the Alexander Terrane in Southeastern Alaska and Northwestern British Columbia. EG <u>103</u>(1): 89-115.

Thorne, K.G., et al., 2008. Characteristics of Mineralization at the Main Zone of the Clarence Stream Gold Deposit, Southwestern New Brunswick, Canada: Evidence for an Intrusion-Related Gold System in the Northern Appalachian Orogen. Explor. Min. Geol. 17(1/2): 13-49.

Tronos, F., et al., 2008. Formation of the Tharsis Massive Sulfide Deposit, Iberian Pyrite Belt: Geological, Lithogeochemical, and Stable Isotope Evidence for Deposition on a Brine Pool. EG <u>103</u>(1): 185-214.

Wagner, T., et al., 2007. Gold upgrading in metamorphosed massive sulfide ore deposits: Direct evidence from laserablation-inductively coupled plasma mass spectrometry analysis of invisible gold. Geology 35(9): 775-780.

Walker, R.J., et al., 2007. Effects of Mother Lode-Type Gold Mineralization on ¹⁸⁷Os/ ¹⁸⁸Os and Platinum Group Element Concentrations in Peridotite, Alleghany District, California. EG 102(6): 1079-1089.

Wang, M., Gao, Y., and Liu, Y., 2008. Progress in the collection of Geogas in China. <u>in</u> Geochemistry: Exploration, Environment, Analysis 8(2): 183-190.

Warren, I., Simmons, S.F., and Mauk, J.L., 2007. Whole-Rock Geochemical Techniques for Evaluating Hydrothermal Alteration Mass Changes, and Compositional Gradients Associated with Epithermal Au-Ag Mineralization. EG 102(5): 923-948.

Watters, S., et al., 2008. Gold Mineralization at the Anomaly A Deposit, Clarence Stream Area, Southwestern New Brunswick: Distal Deposits of a Syn-Deformational Intrusion-Related Gold System: Explor. Min. Geol. <u>17</u>(1/2): 67-84.

Wellmer, F.W., Dalheimer, M., and Wagner, M., 2008. Economic Evaluation in Exploration. 2nd Ed. Springer-Verlag. 250 p.

Wilburn, D.R., 2008. Exploration Review. Min. Eng. $\underline{60}(5)$: 45-57.

Workman, R.K., et al., 2008. Oxygen isotopes in Samoan lavas: Confirmation of continental recycling. Geology <u>36</u> (7): 551-554.

Zhou, J., et al., 2007. Characteristics of the Bumo gold field, Hainan Island, China and gold exploration through soil geochemical surveys. Resource Geol. <u>57</u>(1): 76-89.

Association of Applied Geochemists APPLICATION FOR MEMBERSHIP*




Plea	ase complete only the relevant section for membership. See below fo	r mailing instructions.	/
of A	, wish to app Applied Geochemists. I have read the Code of Ethics of the Association four and abide by them.	ly for election as aMember /Student Member of the Asson and in the event of being elected a Member/ Student Member a	
	EMBER: State Employer and Employee title n actively engaged in scientific or technological work related to applia		ears.
	(employer)	(employment title)	
ST	UDENT MEMBER: Student status must be verified by a Professor o	f your institution or a Fellow of the AAG	
l ce	ertify that the applicant is a full-time student at	in pure or applied (institution)	science
	(Professor/ AAG Fellow Signature)	(Printed Name and Title)	-
Wit	eness my hand this day of, 20	(Signature of applicant)	_
<u>NA</u>	ME AND ADDRESS: PLEASE PRINT (to be completed by applicant	nt)	
Nar	me:	Telephone bus:	
	dress:		
		home:	
		e-mail:	
An	nual Dues :		
All	applications must be accompanied by annual dues . All payments mu	st be in US funds. Select one of the four listed below.	
1	2008 member dues	US\$ 100	
2	2008 student member dues	10	
	- If receipt required, include a self-addressed envelope and add	2	
	- If your check is not drawn from a U.S.A. or Canadian bank, add	15	
		TOTAL	
	rment by check, International Money Order, UNESCO Coupons, Interestable. For credit cards users, minor variations in your billing may		
Тур	e: VISA American Express Master Card Credit card acc	ount number:	
Ex	p. date:Name:	Signature:	

*Application for voting membership (Fellow) requires the sponsorship of three voting members. Request a voting member application from the Association office.

Please note: Your application form will be acknowledged upon receipt. The Admissions Committee reviews all applications and submits recommendations to Council, who will review these recommendations at the next Council Meeting or by correspondence. If no objection is raised the names, addresses and positions of candidates will be listed in the next issue of the AAG Newsletter. If after a minimum of 60 days have elapsed following submission of candidate information to the membership no signed letters objecting to candidates admission are received by the Secretary of the Association from any Member, the Candidate shall be deemed elected, subject to the receipt by the Association of payment of required dues. Send completed application, together with annual dues to:

Association of Applied Geochemists
P.O. Box 26099, 72 Robertson Road, Ottawa, Ontario, CANADA K2H 9R0

NUMBER 140 EXPLORE PAGE 26



- New sensors and data (eg IKONOS, Quickbird, Worldview, GeoEye)
- Earth viewers (eg Google Earth, Microsoft Virtual Earth)
- Field spectrometers, core, downhole, and ore sorting
- Digital elevation model options and applications
- Hyperspectral surveys and scanners
- Mineral and petroleum Exploration
- Environmental and geohazards
- Possible fieldtrip/workshop

To present a talk on any of the topics listed above please submit a title and abstract of less than 300 words to the GRSG Chairman Dan Taranik at dtaranik@angloamerican.co.uk



Newsletter No. 140

SEPTEMBER 2008

Editor: Beth McClenaghan (bmcclena@nrca.gc.ca) Assistant Editor: Elizabeth Bailey (eabailey@usgs.gov) Business Manager:

Sarah Lincoln, (604) 895-4443 (slincoln@barrick.com)

Back Issues contact: Betty Arseneault (office@appliedgeochemists.org)

EXPLORE is published quarterly by the Association of Applied Geochemists, 1330 Ash Court, Thornton, CO 80229 USA.

EXPLORE is a trademark of the Association of Applied Geochemists.

Type and layout of EXPLORE: Vivian Heggie, Heggie Enterprises, Thornton, CO (303) 288-6540; <vjmheggie@comcast.net>

ADVERTISING RATES

Full page (Black & White)	241h x 190w mm	(9.5h x 7.5w in)	US \$ 970
Full page (Color)		,	US\$1165
Half page (Black & White)	241h x 89w mm	(9.5h x 3.5w in)	US \$ 530
or	124h x 190w mm	(4-7/8h x 7.5w in)	
Half page (Color)		· ·	US \$635
Third page (Black & White)	241h x 51w mm	(9.5h x 2w in)	US \$420
or	178h x 89w mm	(7h x 3.5w in)	
Third page (Color)			US \$505
Quarter page (B&W)	124h x 89w mm	(4-7/8h x 3.5w in)	US \$300
or	241h x 41w mm	(9.5h x 1-5/8w in)	
Quarter page (Color)		,	US \$360
Eighth page (Black & White)	60h x 89w mm	(2-3/8h x 3.5w in)	US \$190
Eighth page (Color)			US \$230
Business Card (B&W)	51h x 89w mm	(2h x 3.5w in)	US \$ 50
Business Card (Color)			US \$ 60

Please direct advertising inquiries to:

SARAH A. LINCOLN, BARRICK GOLD CORPORATION STE 700, 1055 WEST GEORGIA ST • VANCOUVER, BC V6E 3P3, CANADA (604) 895-4443 Fax: (604) 684-9831 (slincoln@barrick.com)

EXPLORE Publication Schedule

Quarterly newsletters in March, June, September, December

Deadlines for submission of articles or advertisements:

March newsletter: January 15 June newsletter: April 15 September newsletter: July 15 December newsletter: October 15

Manuscripts should be double-spaced and submitted in digital format using WORD. Photos and figures (colour or black and white) should be submitted as separate digital files and as high resolution jpeg or PDF files. Tables should be submitted as separate digital files in EXCEL format. All scientific/technical articles will be reviewed. All contributions may be edited for clarity or brevity.

Formats for headings, abbreviations, scientific notations, references and figures must follow the Guide to Authors for Geochemistry: Exploration, Environment, Analysis (GEEA) that are posted on the GEEA website at: http://www.geolsoc.org.uk/template.cfm?name=geea_instructions_for_ authors

Submissions should be sent to:

Beth McClenaghan, Geological Survey of Canada, 601 Booth Street, Ottawa, ON, CANADA K1A 0E8 Email: bmcclena@nrcan.gc.ca

THE ASSOCIATION OF APPLIED GEOCHEMISTS

P.O. Box 26099, 72 Robertson Road, Nepean, Ontario K2H 9R0 CANADA • Telephone (613) 828-0199 www.appliedgeochemists.org

OFFICERS

January - December 2008

David R. Cohen, President School of BEES

University of New South Wales Sydney, NSW, 2052 Australia TFI: 612 9385 8084

FAX: 612 9385 1558 email: d.cohen@unsw.edu.au

David B. Smith, Secretary U.S. Geological Survey Box 25046, MS 973 Denver, CO 80225, USA TEL: (303) 236-1849 FAX: (303) 236-3200 email: dsmith@usgs.gov Paul Morris, Vice President Geochemistry Section

Geological Survey of Western Australia 100 Plain Street, East Perth 6004 Western Australia

TEL: 618 9222 3345

email: paul.morris@doir.wa.gov.au

Gwendy E.M. Hall, Treasurer Geological Survey of Canada 601 Booth Street, Room 561 Ottawa, ON K1A 0E8, CANADA TEL: (613) 992-6425 FAX: (613) 992-6425

email: ghall@nrcan.gc.ca

COUNCILLORS

2008-2009 2007-2008 Councillor Emeritus Elizabeth Bailey Sherman Marsh Mark Arundell **David Garnett** Stephen J. Cook David Lentz Jorge Loredo Chris Oates Ray Lett Olle C. Selinus Cliff Stanley

Brazil Northern Europe Germano Melo Jr. J. B. De Smeth Chile Southern Europe **Brian Townley** China

Benedetto De Vivo Southeast Asia Vacant

Southern Africa Theo Davies UK and Republic of Ireland Deirdre M. A. Flight

COMMITTEES

Australian Geoscience Council Representative

Xueqiu Wang

David Garnett

Awards and Medals Committee

Chair: David Kelley William Coker David Lentz Barry W. Smee Brian Townley

Bibliography Committee

L. Graham Closs, Chair Robert G. Garrett Richard K. Glanzman Eric C. Grunsky Peter J. Rogers

Distinguished Lecturer Committee Jeffrey Jaacks, Chair

Election Official

Sherman Marsh

EXPLORE

Beth McClenaghan, Editor email: bmcclena@nrcan.gc.ca Elizabeth Bailey, Assist. Editor Sarah A. Lincoln, Bus. Manager

Geochemistry: Exploration, Environment, Analysis

Gwendy E.M. Hall, Editor-in-Chief e-mail: Ghall@nrcan.gc.ca

Admissions Committee Nigel Radford, Chair Paul Morris

Cliff Stanley

New Membership Committee Robert Jackson, Chair

Regional Councillor Coordinator

Short Course Committee

Colin E. Dunn, Co-Chair Vlad Sopuck, Co-Chair Student Paper Competition

Chair: Dave Cohen Paul Morris Owen Lavin Kurt Kyser

Symposium Committee Paul Morris, Co-Chair Nigel Radford, Co-Chair Eion Cameron Mario Desilets Philippe Freyssinet

Gwendy Hall Virginia McLemore Barry W. Smee Graham F. Taylor

Web Site Committee Robert Eppinger, Chair Webmaster: Andrew Ransom

Betty Arseneault, Business Manager

P.O. Box 26099, 72 Robertson Road, Nepean, ON K2H 9R0 CANADA, TEL: (613) 828-0199 FAX: (613) 828-9288, e-mail: office@appliedgeochemists.org


LIST OF ADVERTISERS

Acme Analytical Laboratories, Ltd	8
Activation Laboratories Ltd	7
ALS/Chemex	9
AMIS	13
Alex Stewart	14
Becquerel Laboratories, IncCanada	11

Intertek Minerals Service	10
ioGlobal	15
Rocklabs	12
SGS	28
Thermo Scientific	3

AVALETCI NOM
ANABISINA
SGS PAGE 28

TARGET SUCCESS



EOCHEMISTRY

Mobile Metal Ion (MMI) geochemistry measures the mobile ions that accumulate in soil above mineralization. MMI is used successfully to reduce the cost of definition drilling programs and locate many deeply buried deposits, with few false positives. MMI geochemistry is now exclusively available at SGS laboratories.

SGS IS THE WORLD'S LEADING INSPECTION, VERIFICATION, **TESTING AND CERTIFICATION COMPANY**

www.sgs.com/geochem ca.min@sgs.com

WHEN YOU NEED TO BE SURE



Paid Advertisement

Newsletter for The Association of Applied Geochemists P.O. Box 48836, 595 Burrard Street, Vancouver, B.C., V7X 1A0, Canada

Please send changes of address to: Association of Applied Geochemists P.O. Box 26099, 72 Robertson Road, Nepean, Ontario, K2H 9R0, Canada · TEL: (613) 828-0199 FAX: (613) 828-9288 e-mail: office@appliedgeochemists.org • http://www.appliedgeochemists.org

> PR SRT STD. U.S. POSTAGE PAID DENVER, CO PERMIT NO. 3550